

Indexed, Abstracted and Cited: [Index Copernicus International \(Poland\)](#), [ISRA Journal Impact Factor](#), [International Impact Factor Services \(IIFS\)](#), [Directory of Research Journals Indexing \(DRJI\)](#), [International Institute of Organized Research and Scientific Indexing Services](#), [Cosmos Science Foundation \(South-East Asia\)](#), [International Innovative Journal Impact Factor](#), [Einstein Institute for Scientific Information {EISI}](#), [Directory of Open Access Scholarly Resources](#), [Science Indexing Library \(UAE\)](#), [Swedish Scientific Publication \(Sweden\)](#), [citefactor.org journals indexing](#), [Directory Indexing of International](#)

World Journal of Biology and Medical Sciences



Published by Society for Advancement of Science®

ISSN 2349-0063 (Online/Electronic)

Volume 7, Issue-1&2, 27-41, January - June, 2020

Journal Impact Factor: 4.197



WJBMS 07/01/09/2020

All rights reserved

A Double Blind Peer Reviewed Journal / Refereed Journal

www.sasjournals.com

wjbmedsc@gmail.com / wjbms.lko@gmail.com

REVIEW ARTICLE

Received: 03/04/2020

Revised: 29/05/2020

Accepted: 30/05/2020

Review on Bovine Schistosomiasis and It's Associated Risk Factors

Banchiayehu Atanew and Asfaw Kelemu

Bahir dar Zuria Woreda Administration, Bahir dar Zuria Woreda Veterinary Clinic, Ethiopia

ABSTRACT

*Schistosome are a trematode, snail-born parasitic of circulatory system in domestic animals and man. Ruminants are usually infected with cercariae by active penetration of the unbroken skin. It is an economically important disease caused by several Schistosoma species and results in economic losses through mortality and morbidity. The geographical distribution of schistosoma species infecting cattle are mainly determined by the distribution of their respective intermediate host snails. The disease affects rural communities particularly those who depend upon irrigation to support their agriculture and drink contaminated water. Effective transmission of schistosomiasis occurs when the schistosome parasites, the aquatic snail hosts and the human or animal definitive hosts meet in space and time in surface water. The pathological changes with the disease are attributed by the adult parasite, cercaria and the eggs of the parasite. Health education, chemotherapy, environmental and biological control as well as provision of clean water have an innumerable role in the control activity of the disease. The use of traditional medicines in the treatment of schistosomiasis are economically important and a growing concern. There are various types of plants like *Phytolacca dodecandra* (Endod) is the most widely used anti molluscicidal properties with minimal side effects used by developing countries and continuous to be used in the modern world. Diagnosis is based primarily on the clinico-pathological picture, parasitological techniques and post mortem findings. The most effective way to control cattle Schistosomiasis in endemic areas is to*

prevent contact between the animals and the parasite. Awareness creation about the disease, destruction of the snail intermediate host population at their breeding sites, either by chemical or biological methods may be considered as a recommendation to prevent the occurrence of bovine schistosomiasis.

Keywords: *Bovine, Schistosomiasis, Epidemiology and Risk factors.*

INTRODUCTION

Ethiopia is localized in the tropical altitudes of Africa and has an extremely diverse topography and a wide range of climatic features and altitudes of agro-ecological zones, which makes the country suitable for different agricultural production systems. This in turn has contributed to the existence of large diversity of farm animal genetic resource in the country [Anon, 2004].

Livestock production constitutes one of the principal means of achieving improved living standards in many regions of the developing world. In sub-Saharan Africa countries livestock plays a crucial role both for the national economy and the livelihood of rural communities. It provides draught power and raw material for industry [ILCA, 2007]. In Ethiopia, livestock contribute about 30 -35 % of agricultural gross domestic product (GDP) and 12-16 % of total GDP [(AAPMDA, 1999].

Ethiopia has one of the largest livestock populations in Africa. According to [CSA, 2015], the country has 57.83 million cattle, 29.33 million sheep, 29.11 million goats, 1.23 million camels, 60.51million poultry, 2.08 million horses, 0.41million mules and 7.88 million donkeys. Livestock play an important role in providing export commodities in the form of live animals and hides and skins [FAO, 2009].

The country has the largest cattle population which helps to improve the livelihoods of the community especially pastoral communities in the country [(CSA, 2014/15]. It serves as productive, financial, social asset, means of investment, important source of cash income and ensuring food security [MOI, 2015].

However, the cattle production and productivity is below the expected potential due to widespread of animal diseases, inadequate and poor quality of animal feed; limited veterinary services; poor selection and breeding and lack of proper technology package. Among the bottle neck challenges of cattle production, trematode parasites are the main once [Yeneneh et al., 2012].

Flukes of ruminants are flat worm (trematodes) parasites living in liver (*Fasciola*) proventriculus (*Paraphistomum*) or blood (*Schistosoma*). The occurrence of the parasites is depending on the suitable ecological conditions for the growth and multiplication of intermediate host (snails) [Urguhart et al., 2003].

Schistosomes are dioecious parasitic flatworms, which live in the vasculature of their mammalian definitive hosts. They are the causative agent of schistosomiasis, a disease of considerable medical and veterinary importance [Rollinson and Southgate, 1993].

Schistosomiasis is snail-borne trematode infection of man, domestic animals and wild animals in different parts of tropical and subtropical countries.

The geographical distribution of bovine schistosomiasis has been determined primarily by the distribution of snail intermediate host particularly *Bulinus* species which are important for the occurrence of disease in bovine species. *Schistosoma bovis* is a species' whose final hosts are bovines, ovines, caprines and whose secondary hosts are small wild ruminants. They are distributed throughout Africa, South West Asia and Mediterranean, Europe [Singh and Chaudhri, 2004].

These parasites cause a significant economic loss which includes death, loss in carcass weight, reduction in milk yield, decline production and productive performances, exposure of animals to other diseases due to secondary complications and cost of treatment expenses [Mogdy et al, 2009]. Moreover; the public significance Schistosomiasis has been reported from different parts of the world including Ethiopia. Human acquire infection through ingestion of metacercariae that are attached to certain aquatic plant and vegetables [Mohammad and Waqtola, 2006].

The major transmitting sites of the disease are small streams all over the highlands of Ethiopia, lakes like Tana, as well as irrigation systems, such as sugar state [Shibru et al., 1989].

The transmission of schistosomiasis takes place only in the place where fresh water snail vector is present and where there is contact between the host and infested water [Okpala, 2004].

Animal and human schistosomiasis is dependent on environmental factors such as moisture, rainfall, temperature, water bodies (stagnant ponds, swamps, streams, rivers, irrigation canals, marshes and dams) and snail intermediate hosts [Li et al., 2012].

Moreover, schistosome infection is closely associated with infested water bodies with traditional grazing and watering systems [Arshad et al., 2011]. These factors tend to be conducive of enzootic schistosomiasis, which is characterized by the high prevalence and significant losses of productivity in ruminant population [Islam, 2011].

Age, sex, breed, season, immunity, anthropogenic/ human, farming system and environmental factors are also playing the role for the prevalence and occurrence of schistosomiasis infection rate in animals. The young parasites cause some damage during migration, but the principal lesions and pathology are associated with passage of the spined eggs through the tissue of the gut lumen of intestine and other organs, and blood sucking habit of the helminthes worm [Soulsby, 1982].

Praziquantel is highly effective for the treatment of bovine schistosomiasis. Treatment of schistosomiasis helps in reversing acute or early chronic disease, preventing complications associated with chronic infection and preventing neuroschistosomiasis. The most effective way to control cattle schistosomiasis in endemic areas is to prevent contact between the animals and the parasite [Richer, 2003]. Therefore the objectives of this review were to explain the etiology, epidemiology, diagnosis, treatment of Bovine Schistosomiasis, and to describe the risk factors, transmission, and economic importance of Bovine Schistosomiasis.

LITERATURE REVIEW

Etiology

Schistosomes are thin, elongated flukes up to 2cm long primarily parasitizing blood vessels of the alimentary and bladder responsible to cause schistosomiasis [Urquhart et al., 1996].

Schistosomiasis or Bilharziasis is a disease caused by trematodes of the genus *Schistosoma* with different species. The taxonomic classification of the organism is presented as kingdom Animalia, Phylum Platyhelminthes, class Trematoda, sub class digenea, Family Schistosomatidae, Genus *Schistosoma* and a total of nineteen [Aiello and Mays 1998]. Different species are described worldwide and they can be differentiated between from their morphological features, lifecycle, host specificity, enzyme and DNA and behavioral [Rollinson and Southgate 1987]. Veterinary important species of *Schistosoma* that cause major impact on domestic animals include *Schistosoma bovis* (ruminants in Africa and Asia), *Schistosoma leiperi* (cattle in Africa), *Schistosoma spindale* (ruminants, horse and pigs in

Asia), *Schistosoma nasale* (ruminants and horse in India), *Schistosoma indicium* (horse, cattle and goats in India buffalo), *Schistosomaja ponicum* (human, and mammals in Asia) and *Schistosoma margrebowiei* (horses, ruminants and elephants in Africa) [Kassaw, 2007].

Morphology

The Schistosome parasites are elongate, unisexual and dimorphic trematodes, which inhabit the blood vessels of their hosts. The female is slender and usually longer than the male and the female is carried in the gutter-like groove, the gynaecophoric canal of the male. The average length of adult male was $(16.7 \pm 1.3\text{mm})$ and it had two suckers (oral and ventral) and distinct gynaecophoric canal and 3-6 testicles situated behind the ventral sucker, and in all specimens showed male holding the threadlike female in gynaecophoric canal. In fact it seems that females cannot mature in absence of the males (Figure 1). As in other flukes the digestive system of *Schistosoma* is blind, i.e. it has no anus but ends in a blind branch, the cecum. *Schistosoma* eggs have species-specific sizes (130-300x40-90 micrometers) and are oval to spindle-shaped, with or without spines [Kahn, 2011]. The Schistosomes are different from most other members of the digenea, the sexes are separate. The term Schistosome or *Schistosoma* means split body and refers to the fact that the males have a ventral groove called gynaecophoric canal [Marquardt and Greive, 2000].




Figure 1. Morphological characteristics of adult *Schistosoma* [CDC, 2012].

Epidemiology

Schistosomiasis is closely associated with large permanent water bodies such as ponds, lakes and marshy pastures. The distribution of the disease has been primarily determined by the distribution of snail intermediate host, particularly *Bulinus* and *Physopsis* species which are important for bovine Schistosomiasis.

Schistosoma species of animals are widely distributed in the tropical and sub-tropical region of the world. These include *Schistosoma bovis* commonly found in cattle, goat and sheep in Africa and Southern Europe, *Schistosoma mattheei* normally parasite of sheep, cattle and other domestic animals in southern parts of Africa, *Schistosoma magrebowiei* in Antelopes in Central Africa and *Schistosoma rodhaini* normally parasitic in wild rodents and in carnivores in Eastern and Central part of Africa. Other species of veterinary importance in Asia include *Schistosoma spindale*, *Schistosoma indicum*, *Schistosoma incognitum*, *Schistosoma nasale* and *Schistosoma leiperi* [Hailu et al., 1998].

Schistosoma bovis has a localized distribution in Ethiopia and has been recorded in slaughter houses in various region of the country. The prevalence of *Schistosoma bovis* has reported from different regions of the country by fecal examination. For example, in Bahir Dar it was 33.8% report by Solomon [Solomon, 2008], 28% in Kemissie by Ameni et al., [2011] in cattle and 15.5% in Bahir Dar by Haile, [2012], in small ruminants were evident.

Intermediate Host

All Schistosomes need snails as their intermediate hosts to complete the asexual phase of their life cycles [Baugh, 1978]. The snail intermediate hosts for bovine Schistosomes belong to the genus *Bulinus*, *Indoplanorbis* and *Planorbis*. An important genus in Africa is *Bulinus*. *Bulinus* are grouped into africanus group, truncatus/tropicus complex, forsicalli group and reticulates group [Bont, 1995]. Intermediate hosts of *Schistosoma mattheei* belong to the *Bulinus africanus* group while those of *Schistosoma bovis* belong to truncatus/tropicus complex as well as to the africanus group and farcically groups. In Ethiopia *Bulinus truncatus*, *Bulinus africanus* and *Bulinus abysinicus* serve as the intermediate hosts of *Schistosoma bovis*. *Schistosoma mattheei* has not been recorded in Ethiopia so far. *Bulinus africanus* and *Bulinus truncates* were identified harboring schistosoma cercariae confirmed by the experimental infection of mice with these cercariae. Other snails identified were *Biomphalari apfeifferi* and *Lymnea natalensis*. *Bulinus* species were identified at all sites [Haile, 1987].

Risk factors for infection

Host related risk factors

Age

Due to a long exposure time because older animals move long distances in search of scarce pastures and water thereby increasing their chances of infection as well as becoming infected at overcrowded watering holes, infection rate increased with the increase of age and peak infection occurred at the maturity of age [Bedarkar et al., 2000]. On the other hand, the very young calf do not graze extensively as the older do, so they get less infection of cercariae unlike adult animals.

Sex

Differences in susceptibility to infection between sexes have been observed by various workers. The observed disparity results in differences in exposure [Magona and Musisi, 2002]. The reason seems to be related to social practice of keeping females under better

management and feeding condition for milk production and breeding whether males are generally let lose to graze freely in pasture and infrequently used for draught purpose and also more stressed. Males are also fed relatively poor diet which increases the susceptibility to parasitic infection [Houdijk and Athana, 2003].

Breed

Local breeds are more affected by schistosomiasis than cross breeds. This difference in prevalence of the disease does not due to the difference in susceptibility but due to the difference in exposure. Cross breeds are mostly kept for dairy or fattening purpose and they are mostly housed and supplementing good feed and clean water which reduce their access to the cercariae. However, the local once are mostly managed extensively to graze freely and get access to infective stage of the parasite [Alemseged, 2010].

Immunity

In older animals, faecal egg counts remain low, tissue egg counts seem to follow the pattern of the faecal egg counts, while worm burden tends to increase with the age of the host [Majid et al., 1980].

This suggests the development of an acquired immunity, which mainly acts through a reduction of the fecundity of the female worm, expressed as reduced faecal, and tissue egg counts, with few effects on worm burden. However, evidence is accumulating that with increasing duration of exposure to continuous challenge, cattle also become less susceptible to reinfection. Examination of naturally infected animals has shown that partial protection against reinfection also occurs, and acquired resistance to schistosomes is of major importance in the regulation of infection intensity in the field.

Reductions in worm burden and egg counts could also be induced in non-immune calves, which received serum from immune donors [Bushara et al., 1994].

A few studies reported on heterologous resistance. Calves previously exposed to infection with the human schistosomes *S. mansoni* and *S. haematobium* were partially protected against *S. mattheei* and *S. bovis*, and it was believed that this type of heterologous resistance might be of considerable importance in protecting cattle from the more serious effects of schistosomiasis [De Bont and Vercruysse, 1997].

Seasonal risk factors

Schistosome infection rate in cattle increases during rainy season, because of the abundance of snails and their rapid multiplication and dispersion. Furthermore, dispersion of fecal matter occurs due to rain splashes. These factors may enhance the infection of snails by miracidia and cercarial contamination to adjacent areas through water. During this time conditions on the lands are suitable for the survival of the intermediate hosts and they become heavily infected with the schistosome larval stages. So, cattle are prone to get the infection of schistosomes [19]. But in dry season infection rate of the schistosome parasite is low because of harsh dry conditions and less chances of infection due to unavailability of snail intermediate hosts as the water sources are scarce in this season.

Management risk factors

Cattle schistosomiasis is dependent on environmental factors such as moisture, rain fall, temperature, presence of water bodies (stagnant, swampy, and marshy) and snail intermediate hosts. Husbandry practice such as grazing system, keeping animals whether they are kept all together and /or separately, feeding and drinking areas [Mersha et al., 2012].

Due to the better management practices and sanitation, infection rate reported to be low in intensive farming system where animals are mainly stall fed proper management practices

and policy change towards urban husbandry can minimize the Schistosomiasis prevalence. They also reported that when cattle are slaughtered through back yard system and consequently the stomach and other intestinal contents including blood and washed materials are dumped into the nearby water bodies' prevalence of the disease also increases [Belayneh and Tadesse, 2012].

In the extensive and semi-intensive system of rearing where animals grazing in the fields have more risks of getting contact with water and subsequently with the infective stage, cercaria. Moreover, increasing cattle mobility through trading and or rental potentially increases the possibility of spreading the disease or infection sources.

Human /Anthropogenic factors

Transmission rates of schistosomiasis increases due to Construction of water schemes to meet the power and agricultural requirements for development [Chitsulo et al., 2000] water development projects are the major factor associated with the increase of schistosomiasis, particularly manmade lakes (hydroelectric power) and irrigation schemes (agriculture), which can lead to shifts in snail vector populations. Many surface irrigation systems in Africa create favorable snail breeding conditions that facilitate the transmission of schistosomiasis [World Health Organization (WHO) 2004].

Factors like the variability within irrigated areas, the canal type, the distance of sites from the canal, the composition and density of aquatic vegetation), the season [World Health Organization (WHO) 2007], specific local conditions such as water stagnation, water depth and shading [Dale and Polasky, 2007] and water flow velocities and the location of breeding sites are affect the presence and density of snails among sites. Low-flow velocities and locations found at the starting point of low-order canals are favorable to get aeration of the water and food availability for snails [Boelee and Madsen, 2006].

Transmission

The infective stage of *Schistosoma* is fully mature, cercariae which are leave in the snail and swim freely in the water, where they remain viable for several hours. Ruminants are usually infected with cercariae by penetration of the skin, although infection may be acquired orally while animals are drinking. During penetration, cercariae develop into schistosomula, which are transported via the lymph and blood to their predilection sites of mesenteric and hepatic veins of the host (except for *S. nasale*, which lives in the nasal veins). Where they feed on blood and produce eggs with a characteristic terminal or lateral spine. Eggs passed in the feces must be deposited in water, hatch and release miracidia, which invade suitable water snails and develop through primary and secondary sporocysts to become cercariae [Fraser et al., 1991]. The prepatent period varies according to the species but is generally 45–70 days. The increased host range of the hybrid parasites and changes in host distribution seen in Africa may have a direct impact on transmission of these schistosomes. Laboratory hybrids have been observed to acquire enhanced characteristics such as infectivity, fecundity, and growth rates.

Life Cycle

Schistosomes are dioecious (unisexual) worms, which are an exception among the trematodes and have an indirect life cycle, while water snail act as an intermediate host belong to the genera *Bullinus* and *planorbis* are required to complete their life cycle [Brown, 1980].

After copulation of male and female schistosomes within the lumen of vein, adult female moves against the venous blood flow in to small venules, when she deposited 200-2000

eggs per day. The ova pass through the wall of blood vessels, and then to adjacent tissues. The ova may be discharged in feces or urine and large amount may be trapped in tissues of the final host. In the water, eggs are hatched and release miracidia. The miracidia then invade suitable water snails and develop into cercariae. When cercariae are fully matured, they leave the snail and invade the final host through the skin or mucous membranes after penetration; cercaria develops into schistosomula, which are transported through the lymph and blood to their predilection sites [Jones et al., 1997].




Figure 2. Life cycle of Schistosoma [Donald et al., 2010].

Pathogenesis and pathology

The fully matured cercariae leave the snail and invade the final host through the skin or mucous membranes. After penetration cercariae develop into schistosomula, which are transported through the lymph and blood to their predilection sites [Jones et al., 1997]. The migration of the eggs may cause mechanical damage and lesions. Moreover, Schistosoma eggs trapped in the tissue elicit granulomatous reaction that is mounted to destruct the eggs. These granulomas consist of several cell types, mainly eosinophils, macrophages and lymphocytes [Olds and Mahamoud, 1980]. In the chronic stages of the disease, the pathology is associated with collagen deposition and fibrosis, resulting in organ damage and dysfunction [Kogulan and Lucey, 2005].

The adult parasites are known to be haematophagous and the worm in the viscera can also cause tissue reaction when it is dead. The adult worms lay eggs in the mesenteric vein and as much as 40-80% eggs laid are retained inside the body and it is the retained eggs and their products that are responsible for most morbidity [Hailu, 1999].

Pathologies of infections are highly associated with the eggs and the intensity of the pathogenic effects depends on the duration of the infection and the number of Schistosomes present. Adult parasites in the vascular system can cause lesions. Cercaria induced skin lesions have been also described [Jones et al., 1997].

Clinical findings

The principal clinical signs are associated with passage of the spined eggs through the tissue of the gut lumen. The young parasites cause some damage during migration, but most of the lesions are due to the irritation produced by the eggs of parasites in the intestine and other organs, and blood sucking habit of the helminthes worm. The helminthes worm may also enter the vesical veins and they may cause hematuria.

Diseased animals shown unthriftiness, anorexia, intermittent diarrhea mixed with blood or mucous, dehydrations unken eyes, pallor of mucus membrane, more over severe emaciation and thirst were also encountered. Systemic reactions were mild however; pale mucous membrane, polypnea and nasal discharge were also seen [Zangana and Aziz, 2012]. Signs associated with chronic hepatic disease may develop when eggs are washed back to the liver by the portal circulation during their penetration of the gut wall. The eggs become lodged in the liver and an intense immunological response results, followed by the formation of a granuloma. A large proportion of the liver may be destroyed and the liver function severely disturbed.

Diagnosis

Diagnosis is based primarily on the clinico-pathological picture, seasonal occurrence, previous history of schistosomiasis in the area or the identification of snail habitats with a history of access to natural water bodies, demonstration of characteristic eggs based on the species in the feces, postmortem examination and hematological tests [Urquhart et al., 1997].

Clinical signs

Weight loss, poor weight gain, diarrhoea, loss of appetite, roughness of the skin, and pale mucous membranes of Cattle are a syndrome developed by *S. bovis*. These signs are usually observed by 6-7weeks after exposure to the infective stage, the cercariae. The severity of these signs increases between the 7th and the 9th week, where the fecal egg counts are highest. However, the clinical signs of the disease are unreliable as other trematodes parasites may produce similar clinical signs.

Postmortem findings

Thousands of visible *S. bovis* adult worms are found in the mesenteric veins during diagnosis at necropsy. Infected livers are diagnosed on the basis of the presence of macroscopic lesions of schistosomiasis visible as white-gray foci under the liver capsule and within the substance of the liver [Hendrix and Robinson, 2006].

Parasitological techniques

Detecting eggs of the parasite in feces or biopsy specimen of the infected animal is the only definitive diagnosis of an active *S. bovis* infection. The eggs of animal *Schistosoma* parasites differ in size and shape; oval in *Schistosoma japonicum* and spindle shape in *Schistosoma bovis* and *Schistosoma mattheei* containing single spine protruding from the shell. The position of the spine on the egg shell is a distinguishing feature; a rudimentary lateral spine in ova of *Schistosoma japonicum* and terminal spines in others sedimentation technique the most commonly used method for detection of fecal egg excretion under field condition. In general when schistosomiasis is suspected, diagnosis is best confirmed by a detailed

postmortem examination which reveal lesion and if mesentery is stretched, the presence of numerous Schistosomes in the veins [Urquhart et al., 1987].

Socio-economic and public health significance of the disease

In addition to the high prevalence, outbreak of the disease and increased susceptibility to other parasitic and bacterial disease, the disease has an economic impact like production losses due to *S. bovis* that result from mortality, delayed growth, partial liver condemnation and poor future reproduction performance and sub clinical infections cause significant losses due to long term effects on animal growth and productive capacity or milk yield and draft power according to [McCauley et al., 1984].

Schistosomiasis is one of the most prevalent parasitic diseases and an important public health problem in many developing countries. Globally, schistosomiasis ranks second among parasitic diseases of socio economic and public health importance and is found in 48 African countries [WHO, 1999. An estimated 779 million people are at risk of schistosomiasis, of whom 106 million (13.6%) live in irrigation schemes or in close proximity to large dam reservoirs and the majority of these infections occur in Sub-Saharan [Touré et al., 2008].

It is prevalent in tropical and subtropical areas especially in poor communities that had low access to safe drinking water and adequate sanitation. It is estimated that at least 90% of those requiring treatment for schistosomiasis live in Africa [WHO, 2012]. It was listed among the 13 diseases classified by World Health Organization as “Neglected Tropical diseases [Hotez et al., 2007].

Treatment

Although schistosomiasis not eradicable, the disease can be prevented and transmission controlled with a single annual dose of praziquantel (Donald, 2005). Main control options are the supply of pure drinking water, diagnosis and treatment of positive cases, management of the environment and control of the intermediate hosts or fresh water snails [Hotez et al., 2007].

Praziquantel is highly effective for the treatment of bovine schistosomiasis. Treatment of schistosomiasis helps in reversing acute or early chronic disease, preventing complications associated with chronic infection and preventing neuroschistosomiasis. The goal of treatment is reduction of egg production via reduction of worm load: this reduces mortality and morbidity. Care has to be exercised in treating clinical cases of schistosomiasis since the dislodgement of the damaged flukes may result in emboli being formed and subsequent occlusion of major mesenteric and portal blood vessels with fatal consequences.

Control and prevention

The most effective way to control cattle schistosomiasis in endemic areas is to prevent contact between the animals and the parasite. Unfortunately, this is not always possible in parts of the world where nomadic conditions of management prevail. Other methods of control include destruction of the snail intermediate host population at transmission sites, either by biological methods or chemical, or their removal by mechanical barriers or snail traps and ecological measures.

Biological control of blood flukes (i.e. using their natural enemies) is so far not feasible. However there are medicinal plants which are affective against external and internal parasites. Ecological measures against the snails that aim to render their habitat unsuitable for survival, such as drainage, removal of water weeds, and increased water flow, are important controlling methods of parasitic trematodes and have water snails as intermediate hosts and frequently are found in the same localities as schistosomes.

From the current available chemical Bayluscide (Niclosamide) and copper sulfate are the choices for molluscicide. In addition to these, a native Ethiopia plant, *phytoplaca dodecandora*, locally known as “endod” is also an effective molluscicide.

In human, the most effective way of controlling Schistosomiasis are the provision of sanitary facilities and piped water since; it reduces human contact with contaminated water.

CONCLUSION AND RECOMMENDATION

Schistosomiasis is a chronic debilitating infection of humans and animals caused by different species of schistosomes cattle schistosomiasis caused by *S. bovis* is one of the major veterinary problems. Occurrence of bovine schistosomiasis is dependent on environmental factors such as moisture, rain fall, temperature, water bodies (stagnant, swampy and marshy) and snail intermediate hosts. Although there has been little recognition of its veterinary significance, cattle schistosomiasis can cause significant loss throughout the world. This is due to the nature of the disease which occurs at subclinical level with long term effect on the animal growth and productivity and increased susceptible to other parasitic and bacterial disease. High prevalence of chronic schistosome infections cause significant losses on a herd basis. Although the economic significance of the disease is mainly attributed to morbidity, mortality, liver condemnation, reduced productivity and poor subsequent reproductive performance.

☐ Therefore, further study on the epidemiology of the disease like malacological and parasitological survey, and mapping high risk areas,

☐ destruction of the snail intermediate host population at their breeding sites, either by chemical or biological methods should be carried out for sound prevention and control of schistosomiasis.

☐ Habitat modification and drainage or increasing of water flow activities should be practiced

☐ Human being after swimming must be washed their bodies with soap and dry with clean towel

☐ Well-planned deworming activities are essential in order to reduce the exposure rate of animals to *Schistosoma* and the impact of schistosome infection in the area.

REFERENCES

Anon, J. (2004). States of Ethiopians animal genetics resources country report. A contribution to the first report on the state of world’s animal genetic resources. Institute of biodiversity conservation (IBC), Addis Ababa Ethiopia, pp: 1-54.

ILCA (2007). Annual Report. Addis Ababa, Ethiopia, pp: 34-38.

Animal, animal products and byproducts Market Development Authority (AAPMDA), 1999. Market Problems and Measures to Be Taken.

CSA, (Central Statistics Authority) (2015). Federal Democratic Republic of Ethiopia, Agricultural sample survey on livestock and livestock production (Private peasant holdings), Statistical Bulletin No. 532, Addis Ababa, Ethiopia, 2, 9-20

FAO, (Food and Agricultural Organizations of the United States), 2009. Livestock sector brief, Ethiopia livestock information, and sector analysis and policy branch. FAO, Rome, Italy, 15-16.

Central Statistical Agency (CSA), 2014/15. Federal Democratic Republic of Ethiopia central statistical agency, agriculture in figure key findings of the 2014/15 agricultural

- sample surveys or all sectors and seasons country summary, FDRECSA, Addis Ababa, Ethiopia.
- Ministry of Information (MOI) (2015).** Export products of Ethiopia. Press release of Ministry of Information, Department of press and audiovisual. Addis Ababa, Ethiopia.
- Yeneneh, A., Kebede, H., Fentahun, T. and Chanie, M. (2012).** Prevalence of cattle flukes infection at Andassa Livestock Research Center in northwest of Ethiopia. *Veterinary Research Forum*, 3 (2):85- 89.
- Urguhart, G., Armour, J., Duncan, A., Dunn, W. and Jennings, F. (2003).** *Veterinary Parasitology* 2nd edn. Black well science. Scotland, 277:117 -120.
- Rollinson, D. and Southgate, V.R. (1993).** The Genus *Schistosoma*. A taxonomic appraisal. In D. Rollinson and A.J.G Simpson (EDs), the biology of Schistosomes from genus to latrines London: Academic press, pp: 1-49.
- Singh, A. and Chaudhri, S. (2004).** Visceral Schistosomiasis of domestic animals in India: humoral immune status of infected cattle, sheep and goat against many poly peptide antigens of *Schistosoma indicum* and *Schistosom spidle*. *Parasite and Immunology* 26:167-175.
- Mogdy, H., Al-Gaabary, A., Salaam, A., Oman, M. and Amperage, G. (2009).** Studies on Paramphistomiasis in ruminants in Kafrelsheikh. *Journal of Veterinary Medicine*, 10: 116-136.
- Mohammad, A. and Waqtola, C. (2006).** Medical Parasitology in Jimma University, Jimma Ethiopia, USAID, pp: 284-300.
- Shibru, T., Getachew, T. and Hailu, B. (1989).** Parasitology. Schistosomiasis in Ethiopia. Addis Ababa. Addis Ababa printing press, pp: 18-26
- Okpala, H.O. (2004).** A survey on the prevalence of schistosomiasis among pupils in Apata and Laranto areas in Jos, Plateau State. Volume 3, issue 1. Federal College of Veterinary and Medical Laboratory Technology, National Veterinary Research Institute, Vom, Plateau State, Nigeria.
- Li, S., Qian, Y., Yang, K. et al. (2012).** Successful outcome of an integrated strategy for the reduction of schistosomiasis transmission in an endemically complex area. *Geosp Hlth*, 6 (2): 215-220.
- Arshad, G., Maqbool, A. and Qamar, M. (2011).** Epidemiology of schistosomiasis in buffaloes under different managemental condition in four districts of Punjab, Pakistan. *J Anim Plnt Sci*, 21(4): 841-843.
- Islam, K.S. (2011).** Schistosomiasis in domestic ruminants in Bangladesh. *Tropical Animal Health and Production*, 7(4): 244-248.
- Soulsby, E. (1982).** Helminthes, arthropods and protozoa of Domesticated animals. London: Bailliere Tindal; pp. 136–160.
- Richer, (2003).** The impact of chemotherapy on morbidity due to schistosomiasis. *Acta trop*: 86 (2-3): 161.
- Urguhart, G.M., J. Armor, J.L. Duncan and F.W. Jenning (1996).** *Veterinary helminthology* Veterinary Parasitology. New York Churchill Livingstone Inc., pp: 114-116.
- Aiello, S.E. and A. Mays (1998).** The Merck veterinary manual. 8th ed. Merck and CO, Inc, White house Station, N.J, U.S.A., pp: 29-31.
- Rollinson, D. and U.R. Southgate (1987).** The Genus *Schistosoma*. A taxonomic appraisal. In: D. Rollinson and A.J.G- Simpson (eds), the biology of Schstosomes from genus to latrines, London Academic press, pp: 1-49.

- Kassaw, A. (2007).** Major Animal Health problems of market oriented livestock Development in Fogera Woreda. DVM Thesis, FVM, Addis Ababa University DebreZeit, Ethiopia.
- Kahn, C.M. (2011).** The Merck veterinary manual. 10th ed. White-house Station, NJ: Merck. and Co., Inc; pp. 273–1036.
- Marquardt, C.R. and Greive, B.R. (2000).** Parasitology and vector biology. 2nd ed. Sandiago. Harcourt academic press, pp: 265-272.
- CDC (2012).** Parasite and health. Schistosomiasis. <http://www.dpd.cdc.gov/dpdx/HTML/Schistosomiasis.htm>.
- Hailu, B., Shibru, T. and Leykun, J. (1998).** Schistosomiasis in Ethiopia and Eritrea. Institute of pathobiology, Addis Ababa University, 3rd edition.
- Solomon, O. (2008).** Observations on the prevalence of *Schistosoma bovis* infection in Bahir Dar area, North Central Ethiopia. DVM Thesis: Faculty of Veterinary medicine, Mekele University, Ethiopia.
- Amine, G., B. Krok and T. Bogale (2001).** Preliminary study on the major bovine trematode infection around Kemissie, Northeastern Ethiopia and treatment trial with Praziquantel. Bulletin of Animal Health and Production in Africa, 49: 62-67.
- Haile, S., Golassa, L. and Mekonnen, Z. (2012).** Prevalence of *Schistosoma mansoni* and effectiveness of Praziquantel in school children in Fincha valley, Ethiopia. Journal of Parasitology and Vector Biology, 4: 25-30.
- Baugh, S.C. (1978).** Cited by Southgate, V.R and Agrawal, M.C. Schistosomiasis in India. Parasitol. Today, 6: 166-168.
- Bont, J.D. (1995).** Cattle schistosomiasis. Host parasite interaction. PhD thesis, Universities, Gent, p. 23-27.
- Haile, S. (1987).** Schistosomiasis in domestic ruminants in Bahir Dar. Proceedings of the first National livestock improvement conference, Addis Ababa, Ethiopia Pp; 122-124.
- Bedarkar, S.N., Narladkar, B.W. and Deshpande, P.D. (2000).** Seasonal prevalence of snail fluke infections in ruminants of Marathwada region. *Journal of Veterinary Parasitology*, 14 (1): 51-54.
- Magona, J.W. and Musisi, G. (2002).** Influence of age, grazing system, season and agro climatic zone on the prevalence and intensity of gastrointestinal strongylosis in Ugandan goats. *Small Ruminants Research*, 44: 187-192.
- Houdijk, J.G.M. and Athana, S.S.A. (2003).** Direct and indirect effects of host nutrition on ruminants gastrointestinal parasites. In: Proceedings of the sixth International symposium on the nutrition of herbivores. Merida Mexico.
- Alemseged, G. (2010).** Prevalence of bovine schistosomiasis in Dembia District Northwest Ethiopia. Dvm thesis, faculty of veterinary medicine. Gondar, Ethiopia.
- Majid, A., A., Marchall, T.F. and Hussein, M.F. (1980).** Observations on cattle schistosome miasis in the Sudan, a study on comparative medicine. I. Epizootical observations on *Schistosoma bovis* in the White Nile Province. *Am J Trop Med Hyg* 29: 435– 441.8.
- Bushara, H.O., Omer, O.H., Malik. K.H.E. and Taylor, M.G. (1994).** The effect of multiple transfers of immune serum on maturing *Schistosoma bovis* infections in calves. *Parasitol Res*; 80: 198–202.
- De Bont, J. and Vercruyse, J. (1997).** The epidemiology and control of cattle schistosomiasis. *Parasitol*. 13:255-262.

- Mersha, C., D. Belay and F. Tewodros (2012).** Prevalence of Cattle schistosomiasis and Associated Risk Factors in Fogera Cattle, South Gondar Zone and Amhara National Regional State, Ethiopia. *Journal of Advanced Veterinary Research*, 2: 53-56.
- Belayneh, L. and G. Tadesse, (2012).** Bovine schistosomiasis: A Threat in Public Health perspective in Bahir Dar Town, Northwest Ethiopia. *Acta Parasitologica Globalis*, 5(1): 1-6.
- Chitsulo, L., Engels, D., Montresor, A. and Savioli, L. (2000).** The global status of schistosomiasis and its control. *Acta Trop.*, 77: 41-51.
- World Health Organization (WHO) (2004).** WHO: Prevention and Control of Schistosomiasis and Soil Transmitted Helminthiasis. WHO Technical Report Series, Geneva.
- World Health Organization (WHO) (2007).** Schistosomiasis. WHO Fact Sheet No 115. Geneva.
- Dale, V.H. and Polasky, S. (2007).** Measures of the effects of agricultural practices on ecosystem services. *Ecol. Econ.* 64:286-296.
- Boelee, E. and Madsen, H. (2006).** Irrigation and schistosomiasis in Africa: Ecological aspects. Colombo, Sri Lanka: International Water Management Institute. IWMI Research Report 99:39.
- Fraser, C.M., Bergeron, J.A., Maya, A. and Susan, E.A. (1991).** The Merck veterinary manual: A handbook of diagnosis, therapy, and disease prevention and control for the veterinarian, 7th edition U.S.A, Merck and Co., Inc. Pp 76-78.
- Brown, D.S. (1980).** Fresh water snails of Africa and their medical importance. London: Taylor and Francis. 482.
- Jones, T.C., Hunt, R.D., King, N.W. (1997).** Veterinary pathology, 6 ed. Lippincott Williams and Wilkins, Maryland, pp: 664-667.
- Donald P. McManus, Darren J. Gray, Yuesheng Li, Zheng Feng, Gail M. Williams, Donald Stewart, Jose Rey Ladino, and Allen G. Ross (2010).** Schistosomiasis in the People's Republic of China: the Era of the Three Gorges Dam.
- Olds, G.R. and Mahamoud, A.A. (1980).** Role of host granulomatous response in murine *Schistosoma mansoni*. Eosinophils-mediated distraction of eggs. The American Society for clinical investigation, 66: 1191-1199.
- Kogulan, P. and Lucey, R.D. (2005).** Schistosomiasis medicine specialties, 6: 1-11.
- Hailu, M. (1999).** Observations on the prevalence and intensity of *Schistosoma bovis* infection in Bahir Dar area, north central Ethiopia. DVM Thesis, Faculty of Veterinary Medicine, Addis Ababa University.
- Zangana, I. and Aziz, K. (2012).** Prevalence and pathological study of Schistosomiasis in sheep in Akra (Dohuk province, North Iraq). *Iraq Journal of veterinary Science* 26:125-130.
- Urquhart, G.M., Armour, J., Duncan, J.L. and Jennings, F.W. (1997).** Veterinary helminthology: Veterinary Parasitology. New York: Churchill Livingstone Inc. Pp. 114-116.
- Hendrix, M.C. and Robinson D.E. (2006).** Diagnostic Parasitology for Veterinary Technicians, 3rd ed. New York: Academic press.
- Urquhart, G.M., Armour, J., Duncan, J.L. and Jennings, F.W. (1987).** Veterinary helminthology: Veterinary Parasitology. New York: Churchill Livingstone Inc., Pp. 114-116.
- McCauley, E.H., A.A. Majid and A. Tayed, (1984).** Economic evaluation of the production impact of bovine schistosomiasis and vaccination in the Sudan. *Preventive Veterinary Medicine*, 2: 735-754.

- WHO. (1999).** Report of the WHO informal consultation of schistosomiasis control. WHO/CDS/SIP/99.2, WHO, Geneva.
- Touré, S., Zhang, Y., Bosqué-Oliva, E., Ky, C., Ouedraogo, A., Koukounari, A., Gabrielli, A.F., Sellin, B., Webster, J.P. and Fenwick, A. (2008).** Two year impact of single praziquantel treatment on infection in the national control program on schistosomiasis in Burkina Faso. *Bulletin of the World Health Organization*.86, 780-787
- WHO (2012).** Schistosomiasis. <http://www.who.int/mediacentre/factsheets/fs115/en/index.html>.
- Hotez, P.J., Molyneux, D.H. and Fenwick, A. (2007).** Control of neglected tropical diseases *The New England Journal of Medicine*, 357: 1018-1027. Doi 10. 1056 /NEJMr064142.
- Donald, G. McNeil, Jr. (2005).** Schistosomiasis (Bilharzia) control and prevention. The Carter Center schistosomiasis control program.

Corresponding author: Asfaw Kelemu (DVM), Bahir dar Zuria Woreda Administration, Bahir dar Zuria Woreda Veterinary Clinic, Ethiopia
Email: asfawkelemu31@gmail.com