

Indexed, Abstracted and Cited: [Index Copernicus International \(Poland\)](#), [Google Scholar](#), [ISRA Journal Impact Factor](#), [International Impact Factor Services \(IIFS\)](#), [Directory of Research Journals Indexing \(DRJI\)](#), [International Institute of Organized Research and Scientific Indexing Services](#), [Cosmos Science Foundation \(South-East Asia\)](#), [International Innovative Journal Impact Factor](#), [Einstein Institute for Scientific Information {EISI}](#), [Directory of Open Access Scholarly Resources](#), [Science Indexing Library \(UAE\)](#), [Swedish Scientific Publication \(Sweden\)](#), [citefactor.org journals indexing](#), [Directory Indexing of International Research Journals](#)

## World Journal of Biology and Medical Sciences

Published by Society for Advancement of Science®

ISSN 2349-0063 (Online/Electronic)

Volume 4, Issue- 2, 57-68, April to June, 2017

Journal Impact Factor: 4.197



WJBMS 04/02/244/2017

All rights reserved

A Double Blind Peer Reviewed Journal / Refereed Journal

[www.sasjournals.com](http://www.sasjournals.com)

[wjbmedsc@gmail.com](mailto:wjbmedsc@gmail.com) / [wjbms.lko@gmail.com](mailto:wjbms.lko@gmail.com)

RESEARCH PAPER

Received: 09/04/2017

Revised: 01/06/2017

Accepted: 02/06/2017

### New Method Identification of *Fusarium* species Isolated from Soil and Plants

Hamid Kheyroodin and \*Sadaf Kheyroodin

Assistant Professor in Semnan University, Iran

\*M.Sc. Student in Urban planning in Mashad Azad University, Iran

#### ABSTRACT

*Fusarium* is a large genus of filamentous fungi, part of a group often referred to as hyphomycetes, widely distributed in soil and associated with plants. Most species are harmless saprobes, and are relatively abundant members of the soil microbial community. Some species produce mycotoxins in cereal crops that can affect human and animal health if they enter the food chain. The main toxins produced by these *Fusarium* species are fumonisins and trichothecenes in this research. The growth of *Fusarium* sp. on PDA Isolated from soil and plants that cultivated on Potato dextrose agar (BAM Media M127) and are common microbiological growth media made from potato infusion, and dextrose. Potato dextrose agar (abbreviated "PDA") is the most widely used medium for growing fungi. We identified some selected *Fusarium* species include, *F. avenaceum*, *F. bubigeum*, *F. culmorum*, *F. graminearum*, *F. langsethiae*, *F. oxysporum*, *F. poae*, *F. sporotrichioides*, *F. tricinctum*, *F. verticillioides*, *F. virguliforme*. The detected species corresponded to *F. proliferatum*, *Fusarium* sp., *F. oxysporum* f. sp. *vanillae* (which was the most numerous and most pathogenic to melon roots and stems and leaves), an undetermined species of *Fusarium*, and *F. proliferatum*, which showed no evidence of producing disease symptoms. **Keywords:** *Fusarium* sp., PDA (Potato dextrose agar) Pathogen.

## INTRODUCTION

The genus includes a number of economically important plant pathogenic species. *Fusarium graminearum* commonly infects barley if there is rain late in the season. It is of economic impact to the malting and brewing industries, as well as feed barley. *Fusarium* contamination in barley can result in head blight, and in extreme contaminations, the barley can appear pink.<sup>[4]</sup> The genome of this wheat and maize pathogen has been sequenced. *F. graminearum* can also cause root rot and seedling blight. The total losses in the US of barley and wheat crops between 1991 and 1996 have been estimated at \$3 billion.<sup>[4]</sup>

*Fusarium oxysporum* f.sp. *cubense* is a fungal plant pathogen that causes Panama disease of banana (*Musa* spp.), also known as fusarium wilt of banana. Panama disease affects a wide range of banana cultivars, which are propagated asexually from offshoots and therefore have very little genetic diversity. Panama disease is one of the most destructive plant diseases of modern times, and caused the commercial disappearance of the once dominant Gros Michel cultivar. A more recent strain also affects the Cavendish cultivars used as a substitute for Gros Michel. It is considered inevitable that this susceptibility will spread globally and commercially wipe out the Cavendish cultivar, for which there are currently no acceptable replacements.

*Fusarium oxysporum* f. sp. *narcissi* causes rotting of the bulbs (basal rot) and yellowing of the leaves of daffodils (*Narcissi*).

Cultivated vanilla has a narrow genetic base because it has been extensively cloned (Minoo *et al.*, 2008), which implies that the species is susceptible to epidemics that greatly impact production *via* a large number of diseases. Among these diseases are stem and root rot caused by *Fusarium oxysporum*, wilting due to *Phytophthora* spp., rust caused by *Calospora vanillae* Masee, rot caused by *Sclerotium* sp., and necrosis or anthracnose produced by *Colletotrichum gloeosporioides* (Penz.) Sacc. (Bhai and Thomas, 2000; Summerell *et al.*, 2003; Guzman, 2004; Talubnak and Soyong, 2010).

Among these diseases, special attention has been given to the rotting of stems and roots caused by *Fusarium*, which was initially suggested as being produced by *F. batatis* Wr. var. *vanillae* (Tucker, 1927). Alconero and Santiago (1969) later indicated that *F. oxysporum* f. sp. *vanillae* was the true cause of this disease.

Studies carried out in Indonesia by Pinaría *et al.* (2010) led to the isolation of 12 species of *Fusarium* that were associated with this disease (*F. decemcellulare*, *F. fujikuroi*, *F. graminearum*, *F. mangiferae*, *F. napiforme*, *F. oxysporum*, *F. polyphialidicum*, *F. proliferatum*, *F. pseudocircinatum*, *F. semitectum*, *F. solani*, and *F. subglutinans*), in which *F. oxysporum* has been proven to be the most common species, followed by *F. solani* and *F. semitectum*. These findings have been confirmed in other vanilla-producing regions, and *Fusarium oxysporum* f. sp. *vanillae* has been identified as the principal agent responsible for the disease (Tombe *et al.*, 1993; Bhai and Kumar, 2008; Pinaría *et al.*, 2010).

Despite stem and root rot constituting most of the disease symptoms that affect vanilla production and having been responsible for huge economic losses over the last decades in countries such as Indonesia (Pinaría *et al.*, 2010), this issue has not been sufficiently studied in Mexico, and little concerning the biology of *Fusarium oxysporum* f. sp. *vanillae* has been investigated.

Thus, the objectives of this research were: (i) to identify the *Fusarium* species isolated from vanilla in Papantla Mexico, (ii) to determine the dominant *Fusarium* species associated with stem and root rot of vanilla, (iii) to amplify and sequence the ITS regions of the species to confirm the morphological identification, and to identify unknown isolates.

## MATERIALS AND METHODS

*Collection and isolation of Fusarium spp.* During 2009 and 2010, vanilla roots, stems, and leaves with symptoms of rot (suggesting the presence of *Fusarium*) were collected from *Vanilla planifolia* plants cultivated on a system of orange trees, as standards, in the region of Papantla, Veracruz (20° 21' 47.57" N and 97° 30' 39.06" W). The pathogens were isolated from the collected material in the laboratory. The samples were washed with running water and disinfected with a 2 % solution of sodium hypochlorite for 1 min and then washed twice with sterile distilled water. Two fragments of tissue (~5 × 5 mm) from the area near the damaged region were removed and placed in Petri dishes containing potato–dextrose–agar culture medium (PDA) (DIBICO) or Czapek dox medium (Bioxon). The colonies that developed were observed again under the microscope to identify isolates that corresponded to the *Fusarium* genus according to Leslie and Summerell (2006). From these isolates, a suspension of spores was obtained, and 200 µL of this suspension was dispersed over Spezieller Nährstoffarmer culture medium (SNA: 1.0 g KH<sub>2</sub>PO<sub>4</sub>; 1.0 g KNO<sub>3</sub>; 0.5 g MgSO<sub>4</sub>·7H<sub>2</sub>O; 0.5 g KCl; 0.5 g glucose, 0.2 g sucrose and 20.0 g agar in 1 L of distilled H<sub>2</sub>O). After 24 h, single germinated conidia were transferred to the same culture medium and incubated for 24 h at 25 ± 1 °C. The single spore cultures were maintained in an active state and used for species identification.

*Morphological and molecular analysis of Fusarium spp.* The morphological identification of the single spore cultures was performed according to Montiel–González *et al.* (2005). To achieve this, a 100–µL aliquot of a suspension of conidia (10<sup>6</sup> mL<sup>-1</sup>) was dispersed over Petri dishes containing SNA medium. After 24 h, three germinated spores were obtained, which were then individually transferred to Petri dishes (100 × 15 mm) containing SNA, PDA or CLA (Carnation Leaf–Piece Agar). Molecular identification of the isolates required total DNA extraction according to Cheng and Jiang (2006). For this process, 1.0 g of mycelia was extracted after ten days of development on PDA at 25 °C. To grind the mycelia, 400 µL of TE buffer (10 mM Tris/HCl, 1 mM EDTA, pH 8.0) and 200 µL of phenol–saturated Tris (2 M, pH 8.0) were added. The solution was then transferred to 1.5–mL tubes and centrifuged at 8,000 g for 5 min to separate the aqueous phase from the organic phase. Subsequently, 160 µL of the supernatant was transferred to a 1.5–mL sterile tube, and 40 µL of TE buffer and 100 µL of chloroform were added.

The solution was centrifuged for 5 min at 13,000 g, and the lysate was purified with chloroform twice until a white interface was achieved. Following this process, 160 µL of the supernatant was transferred to a 1.5–mL tube, 40 µL of TE and 5 µL of RNase (10 mg mL<sup>-1</sup>) were added, and the solution was incubated at 37 °C for 10 min. Finally, 100 µL of chloroform was added, and the solution was mixed by inversion and centrifuged for 5 min at 13,000 g, after which 150 µL of supernatant was transferred to a sterile 1.5–mL tube. The aqueous phase contained the purified DNA.

The integrity of the DNA was evaluated by electrophoresis in 0.8 % agarose gels (TBE 0.5 X) in a horizontal chamber (CONSORT) at 100 V. The gels were stained in 100 mL TBE 1X solution containing 2 µL of ethidium bromide (10 mg mL<sup>-1</sup>) for 20 min and then visualized using a photo–documentation (MicroBis) system.

Internal transcribed spacers 1 and 2 (ITS1 and ITS2) were amplified following the procedure established by Vilgalys and Hester (1990) using the oligonucleotides BMB–CR (5'–GCTTAAGTTCAGCGGGT–3') and LR0 (5'–GCTTAAGTTCAGCGGGT–3'). Amplification was carried out in a reaction mixture of 25 µL containing 10X PCR buffer, 2.5 mM MgCl<sub>2</sub>, 1 U Taq

DNA pol, 0.25 mM dNTPs, 25 pM of each initiator (BMB-CR and LR0), and ~50–100 ng genomic DNA.

The process was performed in a thermal cycler (Mastercycler personal Eppendorf) using the following program: 30 cycles of denaturation at 94 °C for 1 min, annealing at 50 °C for 45 sec and extension at 72 °C for one min followed by a final extension at 72 °C for seven min (Vilgalys and Hester, 1990).

The PCR products were purified using a ChargeSwitch<sup>®</sup> – Pro PCR Clean-up Kit (Invitrogen) following the manufacturer's instructions and then sequenced at the Instituto de Biotecnología, Universidad Nacional Autónoma de México (UNAM) using an Applied Biosystems Sequencer (model 391) and employing the BMB-CR oligonucleotide.

The sequence data were edited using BioEdit version 7.0.5.3 (Hall, 1999) and analyzed via the Basic Local Alignment Search Tool (BLAST) system (GenBank, National Center for Biotechnology Information (NCBI) and FUSARIUM-ID (Geiser *et al.*, 2004).

Phylogenetic and molecular analyses were conducted using MEGA version 5 (Tamura *et al.*, 2011) with the Maximum Parsimony (MP) and Maximum Likelihood methods. The MP tree was obtained using the Close-Neighbor-Interchange algorithm (Nei and Kumar, 2000), and the Maximum Likelihood method was based on the Tamura-Nei model (Tamura and Nei, 1993). The bootstrap consensus tree was inferred from 10,000 replicates (Felsenstein, 1985).

Sequences from *Fusarium* species that were related to vanilla and reported by Pinaría *et al.* (2010) were included as species that, according to BLAST, were related to the ITS sequences; these species included the following: *F. sporotrichioides* ATCC 34914 (AB587025.1), *F. semitectum* IP2239 (AF130380.1), *F. oxysporum* f. sp. *vanillae* DL-1-1 (AY383320.1), *F. oxysporum* f. sp. *vanillae* ML-1-1 (AY387700.1), *F. subglutinans* ATCC 38016 (AY898251.1), *F. nygamai* ATCC 12763 (AY898252.1), *F. oxysporum* ATCC MYA-3970 (FJ614642.1), *F. solani* ATCC MYA-4552 (FJ914886.1), *F. proliferatum* NRRL:13569 (GQ167231.1), *F. verticillioides* (FVU34555), *Gibberella fujikuroi* ATCC 52539 (AY898249.1), *G. moniliformis* (EF453174.1), and *Rhizoctonia solani* ATCC MYA-4649 (HQ263348.1) as an external group. All new sequences were submitted to GenBank (accession numbers KM005079–KM005088).

*Pathogenic variation assays.* On leaves – Healthy vanilla leaves of the same plant were collected and washed with a liquid soap solution and disinfected with 2 % sodium hypochlorite (v/v) for 2 min. The leaves were then rinsed twice with sterile distilled water.

To evaluate the pathogenic variation among the 11 isolates on vanilla leaves, a random design was established and repeated five times, and each test was performed twice. For this process, an ~5-mm-diameter fragment of each colony from each isolate was placed on each leaf and incubated in PDA medium for ten days at 25 ± 1 °C. Healthy vanilla leaves with a fragment of PDA free of microorganisms were employed as a control. The leaves were incubated in a damp chamber (100 % relative humidity) at 25 ± 1 °C for 12 days, and four levels of damage were determined: (1) leaves with no symptoms, (2) leaves with chlorosis, (3) leaves with rot, and (4) leaves showing necrosis or dead leaves.

Lesions that developed on symptomatic plants were excised and rinsed in sterile distilled water before plating on PDA. The plates were incubated as described above, and colonies developing from the leaf sections were purified and morphologically and molecularly identified as described above. Re-isolation of the fungus that was used for inoculation was taken as confirmation of Koch's Postulates.




Fig 1. Show fusarium solani cycle.

On rooted stems – The treatments were established in black bags (10 × 15 cm) with 500 g of soil inoculated with 10 mL of *Fusarium* spp. spore suspension ( $10^6$  spores mL<sup>-1</sup>) after 12 days of growth on the PDA medium at  $26 \pm 1$  °C. An initial suspension of spores was prepared from each strain, and 2 mL of Tween 20 solution (20 %) was added to the Petri dish that contained the mycelium. Each suspension of conidia was emptied into test tubes containing 8 mL of sterile distilled water and mixed using a vortexer (MAXI-MIX II). These tubes provided the basis for the preparation of individual suspensions with a concentration of  $10^6$  mL<sup>-1</sup> spores, as determined by counting in a Neubauer chamber. A control solution was also prepared that contained no fungal inoculate, 2 mL of Tween 20 (20 %) and 8 mL of sterile distilled water.




Fig 2. Show Effect of different of Fusarium Spp on plant.

To evaluate pathogenic variation, rooted stems (~30 cm long) with 2 knots were used. The stems were planted two days after inoculating the soil with the fungus. A few moments before planting, a 1-cm incision was made in the portion of the stem that was in contact with the soil using a sterile blade. The stems were watered three times a week, and the temperature of the greenhouse oscillated between 26–28 °C during the day and between 22–24 °C at night.

The assay was established in randomized blocks, with five repetitions where the 11 isolates of *Fusarium* were evaluated; the test was repeated twice. The level of damage to the stems was evaluated after 60 days, and four levels of damage could be observed: (1) cutting with no symptoms, (2) cutting with chlorosis, (3) cutting with rot, and (4) cutting with necrosis or dead stems. Koch's Postulates were applied to confirm the presence of the inoculated fungus, as described above.

The data obtained from these pathogenic tests (leaves and stems) were analyzed using non-parametric Friedman ANOVA and Kendall's coefficient of concordance analysis. To carry out the non-parametric analysis, a Wilcoxon test was applied. All statistical procedures were performed using the Statistica software (version 7.0).

*Identification of Fusarium spp.* One hundred and eighty-nine isolates of *Fusarium* spp. were collected from the damaged vanilla plant tissues and classified into 11 groups according to the differences observed during their mycelial development (color, form and diameter). Once the isolates were classified, they were observed for variability in the microscopic characteristics of their conidia. Based on the foregoing classifications, the isolates of this study were identified as *F. proliferatum* (group HJAG1), *F. oxysporum* f. sp. *vanillae* (HJAG2, HJAG3, HJAG4, HJAG5, HJAG7, HJAG8, HJAG9, HJAG10 and HJAG11) and *Fusarium* sp. (HJAG6).



www.alamy.com - A1FKFA

**Fig 3. Show Foot root contamination of Fusarium Spp.**

The isolates of *Fusarium oxysporum* produced colonies that were white or violet in color and with areal mycelia and a cottony or fibrous texture. The coloring of the surface of the colonies varied from pink to light or dark violet. The microconidia formed false heads in short monophialides . and abundant uni-bicellular microconidia formed that were oval or elliptical. The macroconidia appeared with 3–5 septa in the form of a canoe, with a large apical cell and base cells in the form of a foot . The chlamydospores appeared to be single and were rarely observed in pairs in cultures after three weeks on CLA medium .

Abundant aerial mycelia were observed in the *Fusarium proliferatum* isolates that were initially white in color and later changed to violet–purple. Monophialides and polyphialides with conidia in chain formation and at the head were observed. The macroconidia had 1–3 septa, a slightly curved apical cell and a scarcely developed base cell . The microconidia were oval in form, with a flat base and no septa. No chlamydospores were observed.

According to the morphological characteristics, it was not possible to identify isolates of the third group at the species level. On culture medium, the presence of white to pale yellow mycelia was reported, and monophialides with no branches on false heads as well as abundant macroconidia with three septa were observed, although some had four or five septa . The apical cell was slightly curved, and the base cell displayed the form of a foot. The presence of abundant oval or elliptically shaped microconidia with no or few septa was also observed. No chlamydospores were observed .

To confirm the morphological identification of the 11 studied strains, molecular analyses were performed. Amplification of the ITS fragment of the isolates generated a product of ~600 bp (GenBank accession numbers KM005079, KM005080, KM005081, KM005082, KM005083, KM005084, KM005085, KM005086, KM005087 and KM005088), which presented a similar score in more than 90 % of the cases with regard to *Gibberella moniliformis*, *F. oxysporum*, *F. proliferatum*, and *Fusarium* sp. when comparing the sequences using BLAST analysis.

The results of the dendrograms obtained by the Maximum Parsimony (MP) algorithm and Maximum Likelihood method were similar, which indicated that the strains of *Fusarium* spp. were grouped into three clades. In the first were eight strains that morphologically corresponded to *F. oxysporum* and specifically grouped with *F. oxysporum* f. sp. *vanillae* DL–1–1 and ML–1–1, as reported by Wang *et al.* (2003). A second group was formed by the *Fusarium* sp. (HJAG6) strain, and the organisms corresponded to the *Gibberella fujikuroi* complex. The third clade grouped the HJAG1 strain with *F. proliferatum* according to its morphological description.

*Pathogenic assays.* On leaves – The results revealed significant differences ( $df = 11$ ,  $P < 0.001$ ) in the pathogenic capacity of the isolates. Two *Fusarium* isolates provided highly pathogenic *F. oxysporum* f. sp. *vanillae* (HJAG2 and HJAG9) that caused a high level of damage (3 = rot and 4 = tissue death). Of these, the HJAG2 isolate had the highest level of pathogenic variation ( $df = 11$ ,  $P = 0.0431$ ). The *Fusarium* sp. HJAG6 and *F. oxysporum* f. sp. *vanillae* HJAG7, HJAG8, HJAG10, and HJAG11 isolates showed moderate levels of pathogenic variation, producing tissue chlorosis and small areas of rot. No significant differences in pathogenic variation were visible among these isolates. The *F. proliferatum* (HJAG1) and *F. oxysporum* f. sp. *vanillae* (HJAG5) isolates did not cause any damage to vanilla leaves and were considered non-pathogenic isolates.

On stems – These results also revealed significant differences ( $df = 12$ ,  $P < 0.001$ ). The *Fusarium proliferatum* HJAG1 isolate did not manifest any disease symptoms on stems or leaves. However, *F. oxysporum* f. sp. *vanillae* HJAG8, which had a moderate pathogenic

effect on leaves, was not considered pathogenic to stems. These isolates did not manifest any significant differences with regard to the control.

Contrary to the results observed for leaves, the HJAG5 (*Fusarium oxysporum* f. sp. *vanillae*) isolate showed moderate pathogenic activity because it provoked symptoms of chlorosis and, on some parts of the stem, tissue rot. The HJAG3 isolate behaved in a similar manner as the same species of fungus, and the HJAG2 and HJAG9 (*F. oxysporum* f. sp. *vanillae*) isolates behaved in a similar manner, with vanilla leaves providing one of the more active pathogenic isolates. The HJAG10 and HJAG11 isolates followed, with both corresponding to *F. oxysporum* f. sp. *vanillae*, while HJAG6 (*Fusarium* sp.), HJAG4 and HJAG7 (*F. oxysporum* f. sp. *vanillae*) increased their level of pathogenic variation on stems, where they induced symptoms of tissue rot, yet these isolates only developed chlorosis on leaves. When plant tissues were inoculated with a mixture of the eleven isolates, a greater level of damage was observed on the inoculated stems, leading to wilting and death.

## DISCUSSION

*Fusarium oxysporum* f. sp. *vanillae* was associated with rot disease on stems and roots of vanilla plants in the present study. These results are in accordance with those of Tombe *et al.* (1993), who morphologically identified 100 isolates of *Fusarium* on damaged vanilla tissues and concluded that all isolates belonged to the *F. oxysporum* species. However, the authors did not mention the presence of other *Fusarium* species, as was reported by Pinaria *et al.* (2010), who isolated twelve species of *Fusarium* from damaged vanilla tissues, with *F. oxysporum*, *F. semitectum*, and *F. solani* being the most predominant.

Similar to the results of Pinaria *et al.* (2010), the majority of isolates in this study corresponded to *Fusarium oxysporum* f. sp. *vanillae*, and only a few corresponded to other species, such as *F. proliferatum*. The morphological characteristics of the colonies of these species on PDA media were similar. The results suggested that the HJAG6 isolate, which could not be identified taxonomically using a combination of morphological and molecular characteristics, was similar to the *Gibberella fujikuroi* complex, *Fusarium subglutinans* and *F. verticillioides*. To achieve greater precision, further molecular studies are required (O'Donnell *et al.*, 1998).

According to these authors, nuclear (EF1 $\alpha$ ) and mitochondrial (mtSSU rDNA) gene sequences are excellent phylogenetic markers for resolving problems within the *Fusarium oxysporum* complex. In addition, the use of the EF1 $\alpha$  gene sequence has demonstrated the capacity to corroborate species identity within the *Gibberella fujikuroi* complex.

The moderate pathogenicity of the HJAG6 strain was underlined due to the chlorosis and stem rot it generated on vanilla leaves. The differences in the results of the pathogenic variation tests exhibited by the various strains of *Fusarium* spp. coincided with those reported by Adame-García *et al.* (2011), who observed strong morphological and pathogenic variability as well as a high degree of polymorphism in esterases among the fungi that cause stem rot in *Vanilla planifolia*.

The HJAG7, HJAG10 and HJAG11 isolates that were also identified as *Fusarium oxysporum* f. sp. *vanillae* showed moderate pathogenic activity because they did not cause the death of any tissue, only chlorosis and, in some cases, small areas of rot. The HJAG5 (*F. oxysporum* f. sp. *vanillae*) isolate exhibited special pathogenic behavior, as it did not develop any disease on the leaves and only had a moderate effect on stems, where it produced symptoms of tissue chlorosis and rot in limited areas. In this regard, no reports of similar work on vanilla have been found.




Fig 4. Show growth of mycelium of fusarium on PDA.




Fig 5. Show growth of mycelium of fusarium oxysporium on PDA.




Fig 6. Mycelial Growth of Pleurotus Spp.

It is important to underline that variable pathogenic activity was found for *Fusarium oxysporum* f. sp. *vanillae* isolates, as they demonstrated differences in the severity of damage to each inoculated tissue and among the vegetative tissues that were used. Given these results, further study is needed to widen the knowledge of the pathogenicity of this species to provide an adequate interpretation of its behavior. According to Summerell *et al.* (2011), this pathogenic variation has not been tested or compared with its vegetative compatibility nor have any molecular analyses with other known members been carried out. Further developments using this type of study will contribute towards obtaining information such as that obtained for *Fusarium oxysporum* f. sp. *cubense*, the agent causing Panama disease. Koenig *et al.* (1997), while studying restriction fragment length polymorphisms (RFLPs) in a global collection of *F. oxysporum* f. sp. *cubense*, discovered that this organism might be closely related to those from other hosts, such as tomato. However, O'Donnell *et al.* (1998) mentioned that fungal agents whose evolutionary origins are independent of other special forms that attack other hosts cause Panama disease.

Another species obtained in this study was *Fusarium proliferatum*, which caused no damage to vanilla tissues (leaves or stems) and is therefore considered to be non-pathogenic. Pinaria *et al.* (2010) indicated that this species is less common (found with a frequency of less than 3 %) and is not pathogenic to vanilla, but it is considered to be a saprophytic species or endophytic colonizer of vanilla in Indonesia. *F. proliferatum* is often isolated as a saprophyte on various parts (roots and stem base) of diseased plants and is reported as pathogenic to crops such as maize (Leslie and Summerell, 2006).

The results obtained from the present study offer valuable information on the association of *Fusarium* species with stem and root rot of vanilla in Mexico. These results, together with studies that are currently in progress, contribute towards a better understanding of the etiology of this disease and open possibilities for future investigations related to its control.

## ACKNOWLEDGEMENTS

We thank from Semnan University in Iran cause assistant laboratory and money for determining this research.

## REFERENCES

- Adame-García J., Trigos-Landa Á.R., Iglesias-Andreu L.G., Flores-Estevez N. and Luna-Rodríguez M. 2011. Isozymic and pathogenic variations of *Fusarium* spp. associated with vanilla stem and root rotting. *Tropical and Subtropical Agroecosystems* **13**:299–306.
- Alconero R. and Santiago A.G. 1969. *Fusaria* pathogenic to vanilla. *Plant Disease Reporter* **53**:854–857.
- Bhai R.S. and Kumar A. 2008. Effect of rhizobacteria on *Phytophthora meadii*, *Fusarium oxysporum* f. sp. *vanillae* and *Colletotrichum vanillae* infecting vanilla. *Journal of Biological Control* **22**:33–41.
- Bhai R.S. and Thomas J. 2000. Phytophthora rot—a new disease of vanilla (*Vanilla planifolia* Andrews) in India. *Journal of Spices and Aromatic Crops* **9**:73–75.
- Cheng H.R. and Jiang N. 2006. Extremely rapid extraction of DNA from bacteria and yeasts. *Biotechnology Letters* **28**:55–59.
- Felsenstein J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* **39**:783–791.

- Geiser D., Jiménez-Gasco M., Kang S., Makalowska I., Veeraraghavan N., Ward T., Zhang N., Kuldau G. and O'Donnell K. 2004. FUSARIUM-ID v. 1.0: A DNA sequence database for identifying *Fusarium*. *European Journal of Plant Pathology* **110**:473–479.
- Guzman C.C. 2004. Vanilla. In: Peter K.V. Ed. *Handbook of Herbs and Spices*, pp. 322–349, Woodhead Publishing, England.
- Hall T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**:95–98.
- Koenig R.L., Ploetz R.C. and Kistler H.C. 1997. *Fusarium oxysporum* f. sp. *cubense* consists of a small number of divergent and globally distributed clonal lineages. *Phytopathology* **87**:915–923.
- Leslie J.F. and Summerell B.A. 2006. *The Fusarium Laboratory Manual*. Blackwell Publications, Iowa.
- Lima D.M.M. 1991. Preservacao de espécies de *Fusarium* sob óleo mineral. *Pesquisa Agropecuária Brasileira* **26**:853–855.
- Minoo D., Jayakumar V.N., Veena S.S., Vimala J., Basha A., Saji K.V., Babu K.N. and Peter K.V. 2008. Genetic variations and interrelationships in *Vanilla planifolia* and few related species as expressed by RAPD polymorphism. *Genetic Resources and Crop Evolution* **55**:459–470.
- Montiel-González L., González-Flores F., Sánchez-García B.M., Guzmán-Rivera S., Gámez-Vázquez F.P., Acosta-Gallegos J.A., Rodríguez-Guerra R., Simpson-Williamson W.J., Cabral-Enciso M. and Mendoza-Elos M. 2005. Especies de *Fusarium* presentes en raíces de frijol (*Phaseolus vulgaris* L.) con daños de pudrición, en cinco estados del centro de México. *Revista Mexicana de Fitopatología* **23**:1–7.
- Nei M. and Kumar S. 2000. *Molecular Evolution and Phylogenetics*. Oxford University Press.
- Nelson P.E., Toussoun T.A. and Marasas W.F.O. 1983. *Fusarium species: An Illustrated Manual for Identification*. Pennsylvania State University Press.
- O'Donnell K., Kistler H.C., Cigelnik E. and Ploetz R.C. 1998. Multiple evolutionary origins of the fungus causing Panama disease of banana: Concordant evidence from nuclear and mitochondrial gene genealogies. *Proceedings of the National Academy of Sciences* **95**:2044–2049.
- Pinaria A.G., Liew E.C.Y. and Burgess L.W. 2010. *Fusarium* species associated with vanilla stem rot in Indonesia. *Australasian Plant Pathology* **39**:176–183.
- Summerell B.A., Leslie J.F., Liew E.C.Y., Laurence M.H., Bullock S., Petrovic T., Bentley A.R., Howard C.G., Peterson S.A., Walsh J.L. and Burgess L.W. 2011. *Fusarium* species associated with plants in Australia. *Fungal Diversity* **46**:1–27.
- Summerell B.A., Salleh B. and Leslie J.F. 2003. A utilitarian approach to *Fusarium* identification. *Plant Disease* **87**:117–128.
- Talubnak C. and Soyong K. 2010. Biological control of vanilla anthracnose using *Emericella nidulans*. *Journal of Agricultural Technology* **6**:47–55.
- Tamura K. and Nei M. 1993. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution* **10**:512–526.
- Tamura K., Peterson D., Peterson N., Stecher G., Nei M. and Kumar S. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* **28**:2731–2739.

- Tombe M., Komoto Y. and Tezuka N. 1993.** Identification and cultural types of *Fusarium* isolates from *Vanilla* in Indonesia. *Industrial Crop Research Journal* **6**:1–5.
- Tucker C.M. 1927.** Vanilla root rot. *Journal of Agricultural Research* **35**:1121–1136.
- Vilgalys R. and Hester M. 1990.** Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**:4238–4246.
- Wang Y.Y., Whitehead M., Zhu Y., Burns A., Hocking T., Ruan X.Y. and Kong Q. 2003.** *Fusarium oxysporum* f. sp. *vanillae* isolate DL–1–1 ITS1, 5.8Sr DNA, ITS 2 sequence. Dept. Plant Pathology, Yunnan Agricultural University.

---

Corresponding author: Dr. Hamid Kheyrodin, Faculty of Desert Science, Semnan University, Iran  
Email: [hkhyrodin@yahoo.com](mailto:hkhyrodin@yahoo.com)