

Indexed, Abstracted and Cited: [Index Copernicus International \(Poland\)](#), [ISRA Journal Impact Factor](#), [International Impact Factor Services \(IIFS\)](#), [Directory of Research Journals Indexing \(DRJI\)](#), [International Institute of Organized Research and Scientific Indexing Services](#), [Cosmos Science Foundation \(South-East Asia\)](#), [International Innovative Journal Impact Factor](#), [Einstein Institute for Scientific Information {EISI}](#), [Directory of Open Access Scholarly Resources](#), [Science Indexing Library \(UAE\)](#), [Swedish Scientific Publication \(Sweden\)](#), [citefactor.org journals indexing](#), [Directory Indexing of International](#)

World Journal of Biology and Medical Sciences



Published by Society for Advancement of Science®

ISSN 2349-0063 (Online/Electronic)

Volume 5, Issue-4, 32-37, October to December, 2018

Journal Impact Factor: 4.197



WJBMS 05/04/1010/2018

All rights reserved

www.sasjournals.com

A Double Blind Peer Reviewed Journal / Refereed Journal

wjbmedsc@gmail.com / wjbms.lko@gmail.com

REVIEW ARTICLE

Received: 08/07/2018

Revised: 17/09/2018

Accepted: 18/09/2018

Typological Bacteriological Quality of the Gharb Water Table-Morocco

N. Said, *M. El harti, **K. EL Bouqdaoui,
***KH. El Kharrim and ***Driss Belghyti

Laboratory Biotechnology, Quality and Environment, Faculty of Sciences,
University Ibn Tofail Kenitra, Morocco

*Engineering Laboratory and Materials, Thermostructural Materials and Polymers Team

**Laboratory of Environmental sciences, Faculty of Science Ain Chok,
Hassan II University, Morocco

***Laboratory of Physics and Microelectronic Materials Automatic and Thermal, Morocco

ABSTRACT

Most waterborne pathogens are distributed world-wide, but outbreaks of some, for instance cholera and hepatitis E, tend to be regional. Dracunculiasis is geographically limited to rural areas in Gharb.

This study was carried out on 60 wells of the Gharb region to know the quality of the water table; the samples were taken in three cities Sidi Kacem, Macheraa Belksiri and Sidi Alal Tazi. Bacteriological standpoint, the results revealed significant contamination of the water table of the Gharb region by faecal coliforms and faecal streptococci, as well as anaerobic spore –réductrices sulphite (ASR). Their densities undergo significant spatial fluctuations; faecal streptococci and anaerobic sulphite -réductrices spores are often present. However, the concentrations of the microbial indicators monitored differ according to the location of the wells relative to the different sources of contamination.

Keywords: Groundwater, Quality, Physicochemical, Gharb and Morocco.

INTRODUCTION

The presence of bacteria in groundwater has been demonstrated by the enumeration of cultured bacteria. Bacteria are adapted to the adverse conditions of the underground environment, particularly low nutrient and oxygen levels [Madsen and Ghiorse, 1993, Balkwill et al., 1997].

The origin of bacteria in various types of groundwater is still under discussion [Mayr, 1974] [Dzeda et al., 1998], most of the groundwater bacteria come from the surface medium by transport during the infiltration of surface water into groundwater [Madsen and Ghiorse, 1993]. According to Van Elsas and Heijnen [1990], optimal survival conditions are essential for the introduction of bacterial microorganisms into the soil, and their transfer to the water table. For Fenchel [2001], the ancestors of native bacteria in the aquatic ecosystems underground would come from anoxygenic microorganisms that have evolved.

Various observations suggest that they are active *in situ* and can play an important geochemical role in the environment and more specifically in the contaminated aquifer [Haack and Bekins, 2000].

In this chapter we aim at microbiological parameters, which can be appreciated by; the micro-organs revivable, total coliforms, fecal contamination indicators (faecal coliforms, faecal streptococci and anaerobic sulphite the gear boxes).

MATERIAL AND METHODS

Study site

The plain of Gharb or Basin-pouring Sebou- is a region located in the North West of Morocco, characterized by a very important agricultural and industrial potential. It is crossed by Sebou.

Study Method

The data presented in this study come from 12 months of water withdrawal from all 60 stations (wells); Samples are taken according to Moroccan standards under the references: NM.ISO 19458; 2009. Once the sample has been taken, the vials are kept in a cooler at 4 °C. This is transported to the laboratory for bacteriological analysis within 8 hours. The geographical coordinates of the wells were obtained using a GPS. The analyzes of the samples of well water are carried out at the level of National Institute Hygiene RABAT « Department of Microbiology and Food Hygiene by the analytical methods of reference. Bacteriological parameters sought as well as the references of the methods used are summarized in the following table 1.

Bacteria are the most diverse living organisms on our planet. They are ubiquitous and abundant; they play a role in the process of the circulation of energy and light in ecosystems. Their distribution on land is limited by only a few factors, namely temperature, pH, and water availability [Hendry, 2006].

The five main groups of water indicator organisms were searched for and counted in the water of the wells and sources surveyed. However, the results of our studies will be presented in the form of curves, representing a temporal variation in the rate of germs research, as well as annual means and standard deviations of microbiological parameters during the study period (February 2016 to January 2017).

RESULTS AND DISCUSSION

Study of bacteriological contamination

Enumeration of culturable microorganisms (GT at 37 ° C)

Revivifiable microorganisms (GT) are used as an indicator of global pollution. It encompasses the set of microorganisms capable of multiplying in air at average temperatures, especially at an optimum growth temperature of between 25 and 40°C. It is a health indicator that allows evaluating the number of CFU (Colony Forming Unit) present in 1 ml of water. These are indicators that reveal the possible presence of bacteriological contamination.

The contamination at the level of all the stations follows the same evolution curve. However, the concentration of GT varies between 277 CFU/1mL and 1397 CFU/1mL for well water (figure 1).

These differences in total germ concentrations at the different stations studied are explained by the contamination of water by bacteria from several sources of pollution (waste accumulation, lost wells, manure and leachate) and which is at the origin of the construction of favorable environments for bacterial development.

The high total germ values found in the groundwater of the study area are similar to those obtained at the groundwater level of Meknes and El hajeb by Belghiti [Belghiti et al., 2013a] and Sadek et al., 2016.

Total coliforms (CT)

Faecal coliforms and streptococci are the most common group of bacteria used for bacteriological examination of water [Poole and Hobson, 1979]. Total coliforms (TC) are Gram-negative bacilli; aerobic and anaerobic optional. They are asporulate and do not have oxidase. They are able to ferment lactose with gas, acids and aldehydes at 37 ° C [WHO, 1972], [Archibald, 2000], [Edberg et al., 1997].

The evolution of contamination by total coliforms, has shown that this variation is in saw, and this contamination is important and can reach a maximum of 2981 CFU/100ml (figure 2), recorded in the wells of Sidi Kacem. It should be noted that certain increases in bacterial densities must be punctual, depending on the nature of the substances infiltrated to the water table, knowing that these substances can inhibit or activate microbial growth (such as P 5 and P 6 wells). Sidi Allal Tazi: located closer to the dump), are all factors related to the variability of the density of bacteria in these wells. The high levels of total coliforms could be explained by the poor protection of these water points, the spreading of manure near the water points and infiltration of surface water. These results corroborate those of Belghiti [2013a]; Gaujous [1995].

Fecal Coliforms (CF)

The presence of fecal coliforms may be an indication of the presence of entero- pathogenic microorganisms [Goodman et al., 1982]. The detection of faecal coliforms in drinking water must be seriously contaminated by fecal matter.

The rate of contamination is accentuated in the city of Sidi Kacem, with peaks in the wells near Oued R'dam was consistent with the study of (Sadek et al., 2016), the recorded maximum is 1636 CFU/100ml (Figure 3), this well largely open and unprotected. We note low levels of faecal coliforms in the waters of the sampled stations in the town of Macheraa Belksiri.

Streptococci Fecal (SF)

The presence of faecal streptococci usually indicates faecal pollution [Castany, 1997]. Faecal coliforms and streptococci are the most common group of bacteria used for bacteriological examination of water [Poole and Hobson, 1979]. They are searched for in water as faecal contamination [Gaujous, 1995]. On the other hand, their survival rate in aquatic ecosystems is sometimes higher than that of faecal coliforms [Goodman et al., 1982].

The mean annual levels of faecal streptococci varied between 73 UFC/100 ml as low average at in Macheraa Station BelKesiri (figure 4). These high values of this bacterium have been mentioned in other regions of Morocco, by other authors such as Aït Boughrous [2007] and El Haissoufi [2011].

Spores of anaerobic bacteria sulfite -réductrices (ASR)

Anaerobic Sulphite the gear boxes (ASR) are obligate anaerobes; Gram positive; sporulated and reduce sulphites to hydrogen sulphides. They can survive longer in water than other indicator microorganisms of faecal pollution (coliforms and streptococci) [Regnault, 1990]. Because of their high resistance, ASR spores can detect old or intermittent faecal pollution [Rodier et al., 2009].

The contamination rate Anaerobic sulphite the gear boxes is more or less constant throughout the course of the year, with a well pronounced peak with an increase of about 452 CFU/L 20m, counted in the P4 well in the Macheraa station Belksiri (figure 5), The values found in most water sources studied exceed the Moroccan drinking water standards (NM 03.7.001. 2006); 0 germ of the gear boxes Anaerobic sulphite in 20 ml sample. These results corroborate those found in other water points studied by El Oualilalami [2014] in the region of Fes.

Typological stations according to bacteriological parameters

Principal Component Analysis performed on a data matrix consisting of 20 rows representing sampled stations and 5 columns representing microbiological parameters showed that total germs at 37 ° C, total coliforms and fecal coliforms contribute negatively to the expression of the F1 axis, which cumulates only 61.04 % of the total variance.

The variance provided by the first two components F1 and F2 from the PCA of all the variables, is satisfactory. Indeed, they allow us to explain 87.13 % of the total variance.

The projection of the stations (described by bacteriological parameters) in terms of the first two factorial axis of the PCA (Figure 6) clearly distinguishes four groups of stations:

- Group G1: this group made up of macheraa wells Belksiri which have an average bacteriological water quality, relatively less contaminated with fecal contamination of germs: Total coliforms, fecal coliforms and anaerobic sulphite gear boxes.
- Group G2: this group only has two wells per three wells which is located next to the landfill in the three cities studied , which have very poor bacteriological water quality, relatively more contaminated with fecal contamination germs: GT at 37 ° C, total coliforms and anaerobic sulphite gear boxes.
- Group G3: this group of wells which have water of medium bacteriological quality.

The results obtained in this study confirm the poor environment of these wells. These results are in agreement with those found by [El Haissoufi et al., 2011] ; EL Ouali Lalami [23] for wells and sources prospected in the Fez region, also with those obtained for the Meknes aquifer and those recorded for the surface water table of Marrakech [Boutin, 1987].

CONCLUSION

The use of well water may nevertheless present potential risks to human health and the environment, although studies in this area are still too rare, particularly in Africa.

On the health front, the real problems with the use of well water are the lack of appropriate treatment and the informal setting that often accompanies this practice. However, the microbiological risks are proven, with a transmission of bacterial diseases.

REFERENCES

- Madsen El., Ghiorse Wc (1993).** Groundwater microbiology: subsurface ecosystem processes. In: Aquatic microbiology: an ecological approach. FORDT.E. (Publisher), Blackwell Scientific Publications, Oxford, 167-213.
- Balkwill, D. L. Reeves, H.R., Drake, G., Reeves, J.Y., Crocker, F.H., Baldwin and K.M. Boone, D.R. (1997).** phylogenetic characterization of bacteria in the subsurface microbial culture collection. FEMS Microbiology Reviews, 20, 201-216.
- Mayr, E. (1974).** Populations, species and evolution. Hermann, Paris, 496 pp.
- Dzeda, B., Kaiser, M. and Mach, S. (1998).** Bacteria and groundwater. Soil and Groundwater Pollution. Civil Engineering Dept., Virginia Tech.
<http://www.cee.vt.edu/programareas/environmental/teach/gwprimer/bacteria.html>.
- Van Elsas J.D. and Heijnen, E.C. (1990).** Methods for the introduction of bacteria into soil. Biol. Fertil. Soils, 10, 127-133.
- Fenchel T. (2001).** Microorganisms (microbes), role of. Encyclical. Biodiversity, 4, 207-219.
- Haack, S. K. and Bekins, B.A. (2000).** Microbial populations in contaminant feathers. Hydrogeology Journal, Vol (8), No. 1, P.63-67.
- Hendry, M. (2006).** Extremophiles : there's more to life. Environmental Chemistry, 3, 75-76.
- Belghiti Chahlaoui, A., Bengoumi, D. and El Moustaine, R. (2013a).** Effect of climatic and anthropic conditions on the variation of the water quality of certain wells in rural environment, case of the water table of the plateau of Meknes (Morocco). ScienceLib Editions Mersenne: 5 (131004) ISSN 2111-4706.
- Belghiti Chahlaoui, A., Bengoumi, D. and El Moustaine, R. (2013b).** Study of physico quality . chemical and bacteriological groundwater of the plio- quaternary aquifer in the region of Meknes (Morocco). Larhyss Journal, ISSN 1112-3680, No. 14, June 2013, pp. 21-36.
- Sadek Sanae, Benel Harkati Fatima, Zhor Fathallah, Dakir Zahra, and Belghyti Driss (2016).** Evaluation of the Total Coliforms Antibiotic Resistance Degree of Hospital Wastewater. World Journal of Biology and Medical Sciences. Volume 3, Issue- 2, 45-50, April to June, 2016
- Poole, N.J. and Hobson, P.N. (1979).** Water pollution and its prevention. Microbial . Ecology . 226-245.
- WHO: World Health Organization (1972).** International standards for drinking water. 3oed, World Health Organization. Geneva.
- Archibald, F. (2000).** The presence of coliform bacteria in Canadian pulp and paper mill water systems - a cause for concern. Water Qual Res J. Canada; 35: 1-22.
- Edberg, S.C., Leclerc, H. and Robertson, J. (1997).** Microbial surface contamination and protection against microbial contamination. II indicators and monitoring parameters for parasites. Critical Reviews in Microbiology; 23: 179-206.
- Gaujous, D. (1995).** Pollution of aquatic environments: Aide memoire. Lavoisier (2nd Edition), Technical Collection and Documentation. Paris (France), 220 p.
- Goodman, R.A., Greenberg, H.B., Mckinley, T.E. and Smith, J.D. (1982).** Norwalk gastroenteritis with water system in rural Georgia community. Archives of Environmental Health 37, p: 358-360.
- Castany, G. (1997).** Hydrogeology, principles and methods. Dunod, Paris, 236p.

- Aït Boughrous, A. (2007).** Biodiversity, ecology and groundwater quality of two arid regions of Morocco: the Tafilalet and the Marrakech region. National Thesis. Cadi Ayyad University, Fac. Sci .Semlalia Marrakech, 207p.
- El Haissoufi, H., Berrada S., Merzouki M., Aabouch M., Bennani L., Benlemlih M., Idir M., Zanibou A., Bennis Y. and El Ouali-Lalami, A. (2011).** Pollution of well water in some districts of the city of Fez, Morocco. *Rev. Microbiol. Ind. San and Environn.* Vol. 5, No. 1, p: 37-68.
- Regnault, J.P. (1990).** General microbiology. Microorganisms of water. Ed. Vigot . Paris. pp : 580-609.
- Rodier, J., Legube, B. Merlet, N. and Coll (2009).** The analysis of water. Natural waters, sewage, sea water. Dunod , 9th ed., 1579 p.
- El Ouali Lalami, A., El-Akhal, F., Berrada, S., Bennani, L., Raiss, N. and S. Maniar (2014).** Evaluation of the hygienic quality of well water and springs by the use of a principal component analysis (PCA): A case study of the Fez region (MOROCCO). *J. Mater. About. Sci.* 5 (S1) (2014) 2333-2344.
- Boutin, C. (1987).** The water of the superficial water tables, a vital but vulnerable natural wealth. The example of rural areas of Morocco. *Rev. Sc. Water*, 6 (3): 357-365.

Corresponding author: N. Said, Laboratory Biotechnology, Quality and Environment, Faculty of Sciences, University Ibn Tofail Kenitra, Morocco
Email: sadek_sanae@hotmail.fr