

World Journal of Biology and Medical Sciences

Published by Society for Advancement of Science®

ISSN 2349-0063 (Online/Electronic)

Volume 1, Issue- 4, 81-88, October-December, 2014



WJBMS 1/3/22/2014

All rights reserved

A Double Blind Peer Reviewed Journal



www.sasjournals.com

wjbmedsc@gmail.com / wjbms.lko@gmail.com

RESEARCH PAPER

Received: 05/08/2014

Revised: 27/10/2014

Accepted: 28/10/2014

Serum Lipid Peroxide and Antioxidant Vitamins (E, C) in Neonates with Respiratory Distress Syndrome

Ahmad S. Elshemy, *Ismail Lotfy Mohamad and

**Mohamad Sayyed Bakheet

Department of Pediatrics, Faculty of Medicine, Al-Azhar University, Assuit, Egypt

*Assiut Children University Hospital, Egypt

**Department of Biochemistry, Faculty of Medicine, Al-Azhar University, Assuit, Egypt

ABSTRACT

Respiratory distress syndrome occurs mostly in premature infants with high risk of oxidative stress, free radicals and other reactive species are constantly generated in vivo and cause oxidative damage to DNA and lipid. Antioxidants delay or inhibit cellular damage mainly through their free radical scavenging property, some of it including vitamins C and E. The study was done on sixty preterm neonates with respiratory distress syndrome (RDS), in addition to twenty apparently healthy full term neonates as normal controls to determine the serum levels of the oxidant lipid peroxide and the antioxidant vitamins E and C. the mean levels of serum lipid peroxide which is the end product of lipid peroxidation were found to be higher with variable degrees of significant differences. The raised serum levels of lipid peroxide in neonates with RDS may be due to activated macrophages releasing highly reactive radicals that may cause local disruption of essential structures, and we found that the mean \pm SD levels of serum vitamin C and E were lower with variable degrees of significant differences.

Key words: Respiratory distress syndrome, vitamin C and E, lipid peroxide free radicals, macrophages.

INTRODUCTION

Respiratory distress syndrome, also known as hyaline membrane disease, occurs mostly in premature infants. The incidence and severity of respiratory distress syndrome are related inversely to the gestational age of the newborn infant (Dizdar et al., 2012). Newborns and particularly preterm infants are at high risk of oxidative stress and they are very susceptible to free radical oxidative damage (Davis and Auten et al., 2010). Oxidative stress refers to a state in which an imbalance between pro-oxidants and antioxidants results in oxidative damage, which has been proposed to play an important role in the pathogenesis of many diseases including respiratory distress syndrome (Da Costa., et al., 2012). Oxidative stress presents numerous opportunities for tissue injury through formation of reactive oxygen/ nitrogen species, free radicals and other reactive species are constantly generated in vivo and cause oxidative damage to DNA and lipid. Evidence for oxidative injury comes from measurements of biochemical markers of lipid per oxidation and protein oxidation. Among important oxidants are lipid peroxide and nitric oxide (Reena et al., 2011). Antioxidants are chemical compounds that can slow or prevent the oxidation of other compounds by elimination of reactive free radical intermediates (Jospeh et al., 2010). These antioxidants delay or inhibit cellular damage mainly through their free radical scavenging property. These low-molecular-weight antioxidants can safely interact with free radicals and terminate the chain reaction before vital molecules are damaged. Some of such antioxidants including glutathione, vitamins C and E,

superoxide dismutase, catalase and uric acid (Lobo et al., 2010).

MATERIALS AND METHODS

The study was done on sixty preterm neonates with respiratory distress syndrome (RDS). In addition to twenty apparently healthy full term neonates as normal controls. From the inpatients of neonatal intensive care units of Alazhar university hospital (assiut) and Assiut university hospital during the period from august 2013 to February 2014. Neonates suffering from respiratory distress due to causes other than respiratory distress syndrome as sepsis, diseases of the respiratory, cardiovascular, or central nervous systems were excluded from the study. Parents of the neonates were informed about the nature of the study and written consent was taken from each. Neonates of our study were classified as Group 1 neonates including 20 apparently healthy full term neonates (G1) as the control group, Group 2 neonates including 30 preterm neonates with RDS and their gestational age from 34 to 36 weeks(G2), and Group3 neonates including 30 preterm neonates with RDS and their gestational age from 30 to 33 weeks(G3). All neonates of the study were subjected to thorough history taking including prenatal, natal and postnatal history. General and neurological examinations were done. Gestational age assessment was performed by the new Ballard score using neuromuscular maturity and physical maturity (Ballard et al., 1991). Chest examination was done to diagnose RDS, patients are considered to have RDS if they have signs of respiratory distress and decreased air entry on auscultation of the chest combined with radiological findings on chest X ray (Evrin et al., 2011) . Radiographic findings in chest X ray include

fine reticular shadowing throughout both lungs with accentuation of the air bronchogram or white lung appearance in severe cases (Falah and Raed .,2008), The following investigations were done to all patients and controls including-Routine investigations including complete blood count (CBC), erythrocyte sedimentation rate ESR, blood glucose level and serum electrolytes Na, K, Ca. Specific investigations including assessment of the serum levels of the oxidant lipid peroxide and the antioxidant vitamins E and C. Serum lipid peroxide is determined by the reaction of thiobarbituric acid with lipid peroxide in acidic medium at temperature of 95c for 30 minutes to form thiobarbituric acid reactive product. The absorbance of the resultant pink product can be measured at 534 nm (ohkawa et al.,1979) and (Meihuipan et al.,2004). Serum vitamin C level is

determined by use of the reagent 2,6-dichlorophenol-indophenol. The absorbance of the sample can be measured at 520nm (Valko et al., 2007). Serum vitamin E level (alpha tocopherol) was measured by use of the reagents absolute ethanol –aldehyde, xylene, ferric chloride solution and standard solution of D-L-alpha tocopherol, the result of the test can be measured at 460nm and 520nm (Valko et al., 2007). Data were expressed as mean \pm SD. SSPS curve was used for analysis of data.

DISCUSSION

Balanced status between the serum levels of oxidants and antioxidants represent the main target of human body metabolism for normal healthy cellular growth and development.

RESULTS

Table 1. Demographic Data of different neonatal groups GI, GII and GIII.

Demographic Data	G I	G II	G III	Significance (P value)		
				I. vs. II	I. vs III	II vs III
Gestational Age (Wks)	38.3 \pm 0.98	34.9 \pm 1.2	31.7 \pm 1.4	NS	<0.05*	NS
Sex (M/F)	9/11	14/16	17/13	NS	NS	NS
Birth Wt (Kg)	3.22 \pm 0.45	2.39 \pm 0.70	1.72 \pm 0.56	< 0.05*	<0.01	<0.05
Length (cm)	49.20 \pm 10.8	47.4 \pm 4.84	43.6 \pm 2.17	NS	< 0.05*	NS
HC (cm)	34.4 \pm 0.66	33.4 \pm 1.51	31.2 \pm 1.84	NS	NS	NS

Wks; weeks. M; male F; female Kg; kilogram cm; centimeter Hc; head circumference. Table (1) Show a variable degree of significant differences regarding to: Gestational age / wks, Weight / kg, length / cm between different studied groups (G1, GII and GIII).

Table 2. APGAR score.

1 Min	1.1±0.31	4.6±0.50	2.90±0.55	<0.01**	<0.001***	<0.05*
5 Min	1.6±0.68	7.8±1.00	6.10±1.05	NS	<0.001***	<0.05*

Table 2 shows a variable degree of significant differences regarding to Apgar score at 1 and 5 minutes between different studied groups (G1, GII and GIII).

Table 3. Mean ± SD of routine investigations in different studied groups.

Variables	G I	G II	G III	Significance (P value)		
				I. vs II	I. vs III	II vs III
Blood glucose level(mg/dl)	79.20±17.38	65.22±13.14	51.90±26.01	<0.01**	<0.001***	<0.01**
ESR (mm/hr)	8.80±2.68	19.06±3.57	18.07±3.66	<0.01**	<0.01**	NS
Complete blood count (CBC)						
Hb(gm/dl)	16.50±1.72	15.17±3.52	14.65±2.79	NS	NS	NS
RBCs (/mm ³)	4.95±0.32	4.63±0.85	4.91±0.95	NS	NS	NS
HCT%	45.12±9.80	43.66±10.87	43.29±10.89	NS	NS	NS
MCV (fL)	99.94±1.42	107.6±6.73	105.5±6.31	<0.05*	<0.05*	NS
MCH (g/dl)	34.68±1.26	34.99±3.56	32.37±3.79	NS	NS	NS
MCHC(g/dl)	34.75±1.16	32.41±2.10	32.76±2.26	NS	NS	NS
platelets (x10 ⁹ /l)	285.50±2.41	203.6±42.84	201.9±54.74	<0.01**	<0.01**	NS
WBCs(/mm ³)	10.42±3.73	9.93±4.03	9.64±3.27	NS	NS	NS
Serum Electrolytes						
Ca(mg/dl)	9.72 ±0.89	8.86 ±1.04	7.93 ±0.95	NS	<0.05*	NS
Na(mmol/L)	141.24±1.09	140.62±9.40	142.41±3.34	NS	NS	NS
K(mmol/L)	4.34±1.03	5.20±0.79	5.30±0.87	NS	NS	NS

Table 3. Show a variable degree of highly significant differences in blood glucose level in between different studied groups. with variable degree of significant differences in blood parameters (ESR , MCV , platelets) in between GI to GII and GI to GIII. Calcium level is significantly lower in GIII compared to G1. But insignificant differences in other remaining elements in between different subjected neonatal groups.

Table 4. Mean \pm SD of Oxidants and Antioxidants of different studied groups. GI, GII and GIII.

Variables	G I	G II	G III	Significance (P value)		
				I. vs II	I. vs III	II vs III
Oxidants						
Serum lipid peroxide (nmol/mL)	9.92 \pm 3.51	15.17 \pm 9.93	19.82 \pm 3.65	<0.01**	<0.001***	<0.05*
Antioxidant Vitamins						
Serum Ascorbic acid (vit.C) (mg/L)	34.90 \pm 16.82	22.05 \pm 6.50	17.30 \pm 5.25	<0.001***	<0.001***	<0.01**
Serum vit.E (mg/L)	4.81 \pm 0.34	2.56 \pm 0.72	1.98 \pm 0.62	<0.01**	<0.001***	<0.05*

Table 4 shows that

Group II compared to Group 1 showed a highly significant difference ($P < 0.01$) for serum lipid peroxide and vit. E and very highly significant difference ($p < 0.001$) in serum ascorbic acid. Group III compared to Group 1 showed very highly significant difference ($P < 0.001$) for serum lipid peroxide and antioxidants vitamins (C and E).

Group III compared to Group II showed significant difference ($p < 0.05$) for serum lipid peroxide and vit. E and a highly significant difference ($p < 0.01$) for serum ascorbic acid.

Newborns, especially preterm, or low birth weight or even full term healthy births are associated with an increased risks from oxidants over burden which places these infants at higher risk of cellular injury but the protective compensatory mechanisms of different human body cells through several antioxidants either individually or cumulatively keep the normal healthy cellular growth and developmental pattern in different human body organs and tissues after birth (Davis and Auten et al., 2010).

The aim of this study was to evaluate the serum levels of lipid peroxide as an oxidant agent in addition to serum levels of vitamin C and E as antioxidants in subjected different neonates with respiratory distress

syndrome (RDS), compared to full term healthy neonates as a control group.

As regard the demographic data of growth parameters of subjected neonates in our study showed lower mean \pm SD levels of birth weight with variable degrees of significant differences in comparing GIII to G II and in comparing both groups to control group and also lower mean \pm SD of length in group II compared to control group. These results were in agreement with (Geoffrey et al. 2005, Saker et al. 2008, Bental et al. ,2011, Jeng et al., 2012).

Regarding to, Apgar score at one / five minutes in different subjected neonatal groups, in the present study, showed a variable degree of significant differences when comparing GII and GIII to control

group, and to each other. Our results were in concordance with previous researches by (Kumar et al., 2012, Dizdar et al., 2012 and Davis and Auten, 2010). These results revealed the impacted and effectiveness of balanced status regarding to oxidants / antioxidants in different studied neonates and maturity state.

Blood glucose levels were significantly lower in G \mathbb{W} compared to G \mathbb{P} and also when comparing both groups to control group. It was stated by Kayıran SM and Gürakan B, 2010 that the pre-term babies had a lower mean blood glucose concentration in the first few postnatal hours compared with term babies.

As regard complete blood picture, we found significant lower counts of blood platelets in RDS subjected neonates (G11 and G111) compared to control group. This finding is confirmed by several studies reported a much higher incidence of thrombocytopenia among sick neonates admitted to the NICU than in the general neonatal population (18%-35% vs < 1%, respectively). (Castle, et al 1986 and Mehta et al., 1980). The highest incidence was found among the smallest and most premature infants, with approximately 70% of neonates born at a weight < 1000 g developing thrombocytopenia at some point during their hospital stay (Christensen, et al., 2006).

The MCV of red blood cells showed higher volume of RBCs in hypoxic RDS neonates but within normal neonatal limited volumes in the present study including GII and GIII compared to control group.

Calcium level is significantly lower in G111 compared to control group.

Other blood investigated parameters showed variable degrees of insignificant differences including hemoglobin Hg, RBCs count, WBCs count, hematocrite value

MCH, MCHC, sodium and potassium between all neonatal groups of our study.

In the present study, the mean levels of serum lipid peroxide which is the end product of lipid peroxidation were found to be higher with variable degrees of significant differences when comparing G111 and G11 to control group and also in G111 compared to G11. The raised serum levels of lipid peroxide in neonates with RDS (G11 and G111) may be due to activated macrophages releasing highly reactive radicals that may cause local disruption of essential structures (membrane lipids, deoxyribonucleic acid and proteins) and hence tissue destruction (Nair et al., 2008). Our results were in concordance with (Kumar, 2008, Evrim et al., 2011 and Da Costa et al., 2012).

In our study, we found that the mean \pm SD levels of serum vitamin C and E were lower with variable degrees of significant differences when comparing G \mathbb{W} and G \mathbb{P} with G1 and with each other.

The role of anti-oxidant vitamins (C and E) in the maintaining of physiological process of fetal growth during pregnancy was reported by Wang et al., 2009.

Brion et al, 2003 confirmed that preterm infants have lower antioxidant vitamins in their serum compared with term controls. Vitamin E is decreased as a result of vitamin C deficiency. This is explained by (Ashok et al., 2008) and (Traber, 2006) who reported that Ascorbate could regenerate tocopherol from the tocophoroxy radical.

REFERENCES

- Ashok Kumar, Rajiv Ranjan, Sriparna Basu, H.D. Khanna and V. Bhargava (2008).** Antioxidant Levels in Cord Blood of Low Birth Weight Newborns, *Indian Pediatric*, Volume 45–July 17.
- Ballard, J.L., Khoury, J.C., Wang, L., Eilers-**

- Walsman, B.L., Lipp, R. (1991).** New Ballard score, expanded to include extremely premature infants. *Journal of Pediatrics* 119 3:417-23.
- Bental, Y., Reichman, B., Shiff, Y., Weisbrod, M., Boyko, V., Lerner-Geva, L. and Mimouni, F.B. (2011).** Collaboration with the Israel Neonatal Network. Impact of maternal diabetes mellitus on mortality and morbidity of preterm infants 24-33 weeks' gestation. *Pediatrics*. Oct; 128 4: e 848-55. doi: 10.1542/peds.2010-3443.
- Brion, L.P., Bell, E.F. and Raghuvver, T.S. (2003).** Vitamin E supplementation for prevention of morbidity and mortality in preterm infants. *Cochrane Database Syst Rev*; CD003665
- Castle, V., Andrew, M., Kelton, J., Giron, D., Johnston, M., and Carter, C. (1986).** Frequency and mechanism of neonatal thrombocytopenia. *J Pediatr* 108 5 Pt 1:749-755.
- Christensen, R.D., Henry, E., Wiedmeier, S.E.(2006).** Thrombocytopenia among extremely low birth weight neonates: data from a multihospital healthcare system. *J Perinatol* 26 6: 348-353.
- Da Costa, Laura, A., García-Bailo, Bibiana Badawi and Alaa El-Soheemy Ahmed (2012).** Genetic determinants of dietary antioxidant status. *Prog Mol Biol Transl Sci*, 108:179-200.
- Davis, J.M. and Auten, R.L. (2010).** Maturation of the antioxidant system and the effects on preterm birth. *Semin Fetal Neonatal Med*; 15 4: 191-5.
- Dizdar, E.A., Sari, F.N., Aydemir, C., Oguz, S.S., Erdeve, O., Uras, N. and Dilmen, U. (2012).** A randomized, controlled trial of poractant alfa versus beractant in the treatment of preterm infants with respiratory distress syndrome. *Am J Perinatol*. Feb; 29 2: 95-100. doi: 10.1055/s-0031-1295648.
- Evrin Alyamac Dizdar, Nurdan Uras, Suna Oguz, Omer Erdeve, Fatma Nur Sari, Cumhuri Aydemir and Ugur Dilmen (2011).** Total antioxidant capacity and total oxidant status after surfactant treatment in preterm infants with respiratory distress syndrome. *Ann Clin Biochem* 48: 462-467. DOI: 10.1258/acb.2011.010285.
- Falah Diab Salih and Raed Haleem Al-Saad (2008).** Chest radiographic finding in neonatal dysnea. *Alkindy col Med J*, Vol 4 (2), 40-44.
- Geoffrey, A. Agrons, M.D., Sherry, E. Courtney, M.D., J. Thomas Stocker, COL, MC, USA and Richard, I. and Markowitz, M.D. (2005).** Lung Disease in Premature Neonates: Radiologic-Pathologic Correlation 1 doi: 10.1148/rg. 254055019. *Radio Graphics*, 25, 1047-1073.
- Jeng, Mei-Jy, Yu-Sheng Lee, Pei-Chen Tsao and Wen-Jue Soong (2012).** Neonatal air leak syndrome and the role of high-frequency ventilation in its prevention. *Journal of the Chinese Medical Association*. Volume 75, Issue 11, Pages 551-559.
- Jospeh, N.M., Monika, S., Alok, M. Alok and R. Shruti (2010).** In vitro and in vivo models for antioxidant activity evaluation: A review. *Int. J. Pharm. Res. Sci* 1; 1-11
- Kayiran, S.M. and Gürakan, B. (2010).** Screening of blood glucose levels in healthy neonates. *Singapore Med J* 2010; 51: 853-5.
- Kumar, A., Ranjan, R., Basu, S., Khanna, H.D. and Bhargava, V. (2008).** Antioxidant levels in cord blood of low birth weight newborns. *Indian Pediatr*; 45 7: 583-5.
- Lobo, V., A. Patil, A. Phtak and N. Chandra (2010).** Free radicals, antioxidants and functional foods. Impact on human health. *Pharmacogn Rev*. Jul-Dec. 4 (8) 118-126.
- Kumar Umesh, Meenakshi Mishra and Veeru Prakash (2012).** Assessment of antioxidant enzymes and free radical scavenging activity of selected medicinal

plants. *Free Radicals and Antioxidants* Vol.2 (3) 68-76.

Mehta, P., Vasa, R., Neumann, L., and Karpatkin, M. (1980). Thrombocytopenia in the high-risk infant. *J. Pediatr.* 97 (5) 791–794.

Meihuipan, K., Arthur, I., Yuan, Li-Zhang, Edward and A. Fisher (2004). Lipid peroxidation and oxidant stress regulate hepatic apolipoprotein B degradation and VLDL production. *J. Clin Invest.* 113 (9) 1227-1278.

Ohkawa, H., Ohish, N. and Yagih, D. (1979). Assay for lipid peroxides in animal tissues by Thiobarbituric acid reaction. *Anal Biochem.* 95: 351-58.

Reena Negi, Deepti Pande, Ashok Kumar, Sriparna Basu, Ranjana S.K. Hanna and Hari D. Khana (2011). In vivo oxidative DNA damage, protein oxidation and lipid peroxidation as a Biomarker of oxidative stress in preterm LBW infants. *Journal of Medical Sciences.* 11: 77-83

Saker, M., N.S. Mokhtari, S.A. Merzouk, H. Merzouk, B. Belarbi and M. Narce (2008). Oxidant and antioxidant status in mother

and their newborns according to birth weight. *Eur. J. Obstet. Gynecol. Reprod. Biol.*, 141: 95-99.

Suhail Mohd, Safia Suhail, Bharat Kumar Gupta, and Vinay Bharatd (2009). Malondialdehyde and Antioxidant Enzymes in Maternal and Cord Blood, and their Correlation in Normotensive and Pre-eclamptic Women. doi: 10.4021 / jocmr 07.1252. *J Clin Med Res.* August; 1 (3) 150–157.

Traber, M.G. Vitamin E. In: Shils M.E., Shike, M., Ross, A.C., Caballero, B., Cousins, R., eds (2006). Modern Nutrition in Health and Disease. 10th ed. Baltimore, MD: Lippincott Williams & Wilkins; 396-411.

Valko M., Leibfritz, D., Moncol, J., Cronin, M.T., Mazur, M. and Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem cell Biol;* 39 (1) 44-84.

Wang, Y.Z., Ren, W.H., Liao, W.Q. and Zhang, G.Y. (2009). Concentrations of antioxidant vitamins in maternal and cord serum and their effect on birth outcomes. *J Nutr Sci Vitaminol Tokyo* ; 55 (1), 1-8.

Corresponding author: Dr. Mohamad Sayyed Bakheet, Department of Biochemistry, Faculty of Medicine, Al-Azhar University, Assuit, Egypt.

Email: sayyed_2006@yahoo.com